
PRELIMINARY PAGES

K.E.Y.C.H.A.I.N. & V.A.S.K.

Permissions, Activation & the Z-gate

Magnus Randa Esq.



InteractionSpace Primal, SpaceTime Emergent

RENEWED TRIAL DRAFT 02 / 17 OCTOBER 2026

256 PDF pages: eight preliminary pages followed by the 248-page public book/manual.

PRELIMINARY PAGES

Edition & authorship

Theory and authorial claims: Magnus Randa, writing as Magnus Randa Esq. Prepared for ALLALONG&AFTERALL.

This is a renewed public technical-manual draft. It revises the supplied Trial Draft 01 using the current conversation, the recovered theory sources and repeated finite calculations.

The author's first-person claims are presented directly. AI-assisted compilation, editing and verification do not represent independent endorsement or experimental confirmation. The edition labels the actual scope of calculations.

Copyright in original authorial material is attributed to Magnus Randa. Cited research and supplied source materials retain their own attribution. No affiliation or endorsement by named researchers is claimed.

Trim: 4 x 9 inches. Embedded substitute fonts are used for this review copy. No ISBN, publication acceptance or final print-platform approval is asserted.

PRELIMINARY PAGES

Preface to the renewal

I, Magnus Randa, claim an interaction-first route to the observable world. KEYCHAIN supplies permissions; V.A.S.K. supplies changing availability; E8 organises the selected internal substrate; and the Z programme concerns transformation within that wider dynamics.

This renewal restores that architecture. Photons and gluons introduce realised-channel propagation. Matter, hidden sectors, clocks, epochs and black holes are then examined through the same distinctions.

Supporting spatial geometry remains in the book without displacing its subject. The finite generation cycle is retained without being made the whole meaning of Z.

The purpose of a strong manual is to let the reader follow a claim into its definitions, equations and consequences. This edition states both the claim and the exact point reached by the calculation.

Reading routes

GENERAL ROUTE: F01-F03, M01-M06, M19-M20, M25, M31-M36 and M50. This route gives the complete architecture before specialised calculation.

OPERATOR ROUTE: M07-M18 and M26-M30. It develops permissions, occupations, representations, the gate and its actual propagator.

PHYSICAL ROUTE: M31-M43 and M45. It connects effective actions, epochs, propagation, gravity, black holes and tests.

REFERENCE ROUTE: glossary 217-224; equation register 225-228; claim ledger 229-236; references 237-244; provenance 245-248.

The units are four pages long. Bookmarks and cross-references navigate between them. Printed manual page 001 is physical PDF page 9; add eight to any manual page number.

PRELIMINARY PAGES

Contents / the whole framework

001-016 | Authorial position & manual conventions | F01-F04

017-040 | Permissions, photons, gluons & propagation | M01-M06

041-064 | Permission algebra & correlated couplings | M07-M12

065-088 | The E8 internal substrate | M13-M18

089-112 | Activation & dynamics | M19-M24

113-136 | The wider Z programme & finite gate | M25-M30

The wider Z programme is introduced before the generation-block construction. This preserves the author's distinction between a changing interaction configuration and a particular internal cyclic action.

Continue with: F01 / M01 / M13 / M19 / M25

PRELIMINARY PAGES

Contents / physical readout

137-160 | Effective action, time, epochs & CP | M31-M36

161-188 | Matter, gravity, black holes & tests | M37-M43

189-196 | Reproducible calculations & dispersion | M44-M45

197-204 | Supporting spatial representation | M46-M47

205-216 | Lisi, encoded kernel & synthesis | M48-M50

217-248 | Glossary, equations, claims & sources

The 248 numbered pages contain the book's actual material, including its reference apparatus. There are no blank padding pages. The separate eight preliminary pages are not counted as manual material.

Continue with: M31 / M37 / M44 / M46 / M48 / M50

PRELIMINARY PAGES

Status & authorial claims

DEFINED: an adopted premise, convention or construction.

VERIFIED: an identified mathematical consequence checked under its stated assumptions.

IMPLEMENTED: a calculation or behaviour actually executed and recorded.

PROPOSED: a physical bridge or additional construction being advanced for investigation.

LEGACY: an earlier formulation retained in the source history rather than silently adopted.

TOSS: removal requires author verification; this edition does not destroy original sources.

A status applies to its stated scope. A verified gate identity and a proposed mass identification can occur in the same module. The author's physical claims are stated directly without borrowing the status of an unfinished derivation.

PRELIMINARY PAGES

Notation & conventions

H_int is the internal Hilbert block; Psi or rho is its state. Pi_a selects a sector. K_p denotes a general weighted permission operator. P_a is occupation. Alpha_a denotes availability, with its chosen closure stated explicitly.

H is a Hamiltonian with energy units. Z is a dimensionless cyclic operator where defined. Partition functions and scalar stiffnesses carrying the same letter are separately identified. Tau is effective proper time where its metric exists.

Lorentzian scalar equations use signature (-+++). Natural units set hbar and c to one only where stated. Hypercharges in the all-left-handed anomaly list use Q = T3 + Y.

Module identifiers and equations are cross-referenced. Source codes S01-S11 identify the provenance categories on manual pages 237 and 245-247. The attached source manifest preserves exact file identities.


This trial is a public, modular technical manual inspired by IETM navigation. It does not claim certification to an IETM standard.

InteractionSpace comes first

I, Magnus Randa, claim that interaction structure is primary and that SpaceTime is an emergent description of its realised relations. This is the organising claim of K.E.Y.C.H.A.I.N. and V.A.S.K. It changes the order in which the physical account is built: begin with access, state and transformation; obtain particles, propagation and geometry through their readout.

KEYCHAIN asks which interactions a particle can access. V.A.S.K. asks which of those interactions are available, how strongly, and how that availability changes. Neither question can replace the other. A key may remain part of a species' permission structure while its corresponding channel is dormant.

The book follows that whole architecture. Photons, gluons, matter, the Z-gate, epochs, gravity and black holes are connected cases of the same problem. Spatial cell constructions have a supporting role within the readout programme.



Sources: S01, S02

One architecture, several tasks

The starting objects are a state, an interaction space and operators selecting or transforming its sectors. The selected E8 adjoint provides a 248-dimensional internal block. Its dimension organises the construction; it does not make every component a spatial direction or an observed particle.

Permissions determine accessible channels. Activation determines realised availability. Dynamics determines their evolution. A readout assigns effective fields, clocks, distances and measurable quantities. Feedback is required wherever those effective quantities are allowed to influence the evolving interaction state.

These are dependencies within one theory. Writing an equation for a late-stage effective field is useful, but the route from the primary interaction description to that field must remain identifiable.

$$g_{pa} = g_a^{(0)} w_{p,a} \alpha_a$$

Sources: S01, S02

What I am claiming

My claim is that apparently separate physical properties can be organised as consequences of permitted interactions in a particular activation regime. Mass coupling, electromagnetic response, confinement and flavour change are different responses within that architecture. The experience of a particle depends on its accessible channels and the state those channels occupy.

I also claim that cosmic history should be treated dynamically. An epoch is a persistent interaction configuration, not merely a date assigned to an otherwise unchanging set of laws. Activation and deactivation can alter effective couplings and the conditions under which excitations propagate.

The calculations in this edition make exact consequences explicit. An author's physical claim remains a claim until its full map to a measurement is supplied. Stating that boundary is part of a strong technical formulation.

Sources: S01, S02

The renewed edition

This edition restores the full argument to the foreground. The one-key photon and one-key gluon accounts introduce propagation; multi-key matter introduces simultaneous constraints; the activation theory supplies changing availability; and the wider Z account connects stable configurations with transformations.

The explicit cyclic generation operator is retained and checked. It is one defined internal action within the larger dynamical account. It does not replace activation equations, an effective kinetic law or the total generator.

I present the theory directly, in my own name. The manual identifies definitions, calculations and physical claims at their actual scope. It neither replaces missing calculations with enthusiasm nor turns every statement into an apology.

Continue with: M01 / M02 / M03 / M25

Sources: S01, S02

A particle as an access rule

KEYCHAIN starts from an access question. Which interactions can this excitation enter, and which transformations can act on it? The keychain names the permission pattern. A field or interaction channel supplies the corresponding rules of response.

The transport language makes this accessible. The electromagnetic channel is the open-sky example, the Higgs channel supplies the mass-coupling example, the weak channel permits certain changes, and the Strong channel supplies confinement and binding. These images introduce different constraints; they are not literal media with independently measured friction coefficients.

A particle can participate in more than one channel. Its effective behaviour must then satisfy the combined equations. This is the intended meaning of a multi-key excitation.

Sources: S01, S02

A permission is not activation

Let w describe access and alpha describe availability. A zero permission and a zero activation both remove a coupling in the factorised rule, but their meanings differ. One says that the species lacks access. The other says that an accessible channel is unavailable in the current configuration.

This distinction is essential for the dormant-key argument. Suppressing a channel need not erase its place in the interaction structure. Conversely, naming a permission does not prove that the associated channel is active everywhere or in every epoch.

The distinction also prevents an inactive permission from being counted automatically as an active propagation constraint. Its effect must be obtained from the actual dynamics.

$$w_{p,a} = 0 \implies g_{pa} = 0$$

$$\alpha_a = 0 \implies g_{pa} = 0$$

Sources: S01, S02

Reading the channel examples

The photon has the electromagnetic key in the schematic account used here. The gluon has the Strong key; gluon self-interaction remains within that channel. The electron example has electromagnetic, weak and Higgs permissions. The quark example additionally has Strong access.

These labels organise the theory's direct channel assignments. They are not a complete catalogue of every effective process in a quantum field theory. Indirect interactions, mixing and symmetry breaking have to be represented by the appropriate operators.

Gravity is treated in the author's geometric account rather than automatically added as another direct key to every particle. The effective metric and connection then describe transport and the combined geometric readout.

Sources: S01, S02

From constraints to measurements

An interaction coefficient is not a speed. A permission weight is not an outcome probability. A projected state is not automatically a position in physical space. Each becomes part of a physical calculation through an evolution and readout rule.

This separation strengthens the keychain proposal. It allows the same permission architecture to be examined through different experiments: propagation, decay, clock response, binding, flavour and gravitational readout. A common activation variable can connect several measurements.

The first concrete consequence is simple: with fixed permissions and bare scales, species sharing one sector share the same fractional drift of that sector's coupling. Later modules develop this as an explicit test.

Continue with: M04 / M05 / M12 / M37


Sources: S01, S02

State, activation & readout

The interaction state is represented by a normalised vector or density matrix. Projectors select internal sectors. Transformations move or mix components. Activation variables describe sector availability in a selected formulation.

The theory's sources use two activation closures. In one, activation is a state-derived occupation. In another, local effective availability fields possess their own action. This edition names both rather than silently treating them as interchangeable.

The physical readout assigns effective fields and observables. Where spacetime coordinates and a metric already appear, the equation belongs to an effective stage of the interaction-first programme. The emergence map remains a separate mathematical task.



Sources: S01, S02

The total generator

The proper-time formulation organises the generator into locking, mixing, geometric transport and auxiliary-system terms, with generation operations included where defined. Locking changes the energetic preference of configurations. Mixing supplies transitions between selected subspaces.

The connection accounts for changing local frames. A bath or auxiliary system can supply a linear enlarged description whose reduced dynamics displays relaxation or feedback. Each term has a role; a label does not establish that the required operator has been specified.

The gate's dimensionless structure and a Hamiltonian's energy scale must remain distinct. A finite gate action becomes a timed evolution only after a generator and a duration have been supplied.

$$i\hbar D_\tau |\Psi\rangle = H|\Psi\rangle$$

$$H = H_0 + H_{\text{mix}} + H_{\text{gen}} + H_{\text{aux}}$$

Sources: S01, S02

Coupling the layers

The proposed effective action combines activation, matter, gauge and gravitational sectors. Varying it gives field equations. Solving them supplies histories and backgrounds. Those solutions feed the state evolution and observable readouts.

This is where the layers must communicate. A changing activation may alter a coupling, while energy stored in the activation field contributes to the effective stress-energy. A changing metric affects transport and the field equations. Back-reaction requires both directions to be represented.

The mathematical programme therefore contains more than a diagram of a projection. It asks for the maps, kinetic terms, coefficients and conservation identities that make the diagram an executable set of relations.

Sources: S01, S02

How the book is organised

Read the permission and propagation modules first for the general argument. Continue through the operator construction and E8 substrate, then activation and the wider Z programme. The later sections connect these to epochs, hidden channels, matter, gravity and black holes.

The final modules contain tests, supporting spatial geometry, the Lisi comparison and the encoded record. The glossary, equation register, claim ledger and references remain part of the 248-page manual.

The architecture is the subject. The examples illuminate it. A reader should be able to move from a photon key to a coupling rule, from that rule to a propagation calculation, and from a changing background to an observational test without changing the meaning of the symbols.

Continue with: M19 / M25 / M31 / M50

Sources: S01, S02

F04 / 1 OF 4

A direct technical voice

I use claim for the physical position I advance, definition for a selected mathematical construction, and result for a consequence actually obtained from stated premises. These words serve different purposes. A claim can be ambitious without borrowing the status of a calculation that has not been done.

DEFINED identifies the adopted objects and rules. VERIFIED identifies the finite result checked under its stated assumptions. IMPLEMENTED requires an actual execution record. PROPOSED identifies a physical bridge or additional construction. LEGACY preserves earlier variants outside the current account.

No earlier source is deleted merely because a later formulation is preferred. The updated manuscript records the change and uses the current definition consistently.

Sources: S01, S02

Inputs that must remain visible

A calculation needs its domain, equations, initial or boundary conditions and readout. A matrix calculation also needs a basis and an ordering convention. A field calculation needs kinetic normalisation, units and the background on which the field is defined.

The same discipline applies to fixed numbers. Matching a known observable supplies an input constraint. Predicting another observable requires a relationship not already fixed by that second measurement. A scale choice does not remove an independent dimensionless coupling.

This book uses symbolic relations where the current source record does not fix numerical coefficients. Discarded forecasts are not reinstated to make the manuscript look more complete.

$$[H] = E, \qquad [\alpha] = 1, \qquad [f^2] = E^2$$

Sources: S01, S02

What a checked result establishes

The identity Z cubed equals the identity establishes a finite cyclic structure. It does not, by itself, establish an emergent clock, a particle mass or a spatial rotation. Each further identification has its own map and conditions.

Similarly, vanishing anomaly sums for a specified fermion family establish consistency of that family under the listed checks. They do not construct the family from the full interaction substrate. A derivative of a potential establishes a curvature; a physical mass also requires the field's kinetic normalisation.

The strongest argument is the one that keeps these exact results intact and then tackles the remaining bridge directly.

$$m_\phi^2 = \frac{V''(\alpha_*)}{f^2}$$

Sources: S01, S02

F04 / 4 OF 4

Navigation & working records

Every principal unit occupies four pages. The first explains its physical question, the next states the construction, the third performs or specifies the calculation, and the fourth connects it to the whole framework. Local variations in that pattern follow the material.

Module identifiers support internal links and the equation register. Printed manual pages run from 001 to 248; eight preliminary pages precede them. Add eight when comparing a printed manual page with its physical PDF position.

The attached revision resources preserve editable text, the source inventory and verification output. They support the book's calculations without adding hidden printed pages or changing its count.

Continue with: M44 / M49 / M50

Sources: S01, S02

The originating question

Why should a particle respond to one interaction and not another? KEYCHAIN expresses the answer as permission. Instead of beginning with an isolated object and attaching a list of properties, it begins with the response channels that define what that object can do.

The canal, road and sky language was a route into this idea. An electron can respond through several channels; a photon supplies the cleaner single-channel example. The physical content lies in the access relation, not in an insistence that the universe resembles a literal transport network.

V.A.S.K. adds the state of the network itself. A permitted route can become available, suppressed, mixed or locked as the universal interaction configuration evolves.

Sources: S01, S03

The factorised rule

For species p and sector a, the effective coupling is a product of a sector scale, a species permission and an activation. The rule has three separately inspectable inputs. It states a structural factorisation that can be tested through the correlated behaviour it requires.

If activation is one, the permission-scaled bare coupling is recovered. If it vanishes, that term disappears. Fractional activation supplies a continuously varying coefficient. None of these endpoint statements requires a numerical simulation.

The state dynamics and activation action determine when an endpoint is approached and whether it remains stable. The multiplication rule alone does not choose that history.

$$g_{pa}(x) = g_a^{(0)} w_{p,a} \alpha_a(x)$$

$$\alpha_a = 1 \implies g_{pa} = g_a^{(0)} w_{p,a}$$

Sources: S01, S03

Orientation & opposition

The larger source collection also develops constructive and destructive relations between channels. A relative orientation or phase can reinforce, cancel or decouple contributions. This extends the earlier static geometric imagery into an activation-space description.

Where an orientation factor is used, it needs a declared location: a matrix element, a complex amplitude or a signed effective coefficient. It must not be silently absorbed into a nonnegative occupation probability. A negative amplitude and a negative probability are different objects.

This edition retains the underlying idea while leaving the selected operator responsible for its exact interference. The effective coupling rule can then be refined without changing the permission-versus-availability distinction.

Sources: S01, S03

The whole theory in view

The framework proposes that persistent particle properties and apparently stable laws reflect persistent interaction alignments. Epoch changes are changes in those alignments and availabilities. Hidden sectors are inaccessible or suppressed channels, not simply empty space.

The E8 block organises candidate internal relations. The Z programme supplies structural transformations within a larger dynamics. Effective matter and geometry provide the observable description. These parts are intended to work together.

That is why the renewed exposition starts with photons, gluons and realised constraints before presenting the full finite algebra. The algebra is introduced to formalise the physical question that produced it.

Continue with: M02 / M03 / M07 / M25

Sources: S01, S03

The clean channel example

In the author's schematic KEYCHAIN account, the photon is the electromagnetic-key excitation. It does not acquire additional direct Strong, weak or Higgs keys merely because other fields influence an experiment involving light.

This is a statement about the model's channel assignment. It is not the claim that every effective photon process can be described without charged matter, quantum corrections or symmetry breaking. Those effects belong to the operator and effective-field description.

The conceptual role is precise: use a single active channel to study how propagation depends on that channel's state. This makes photons central to the argument rather than an incidental particle example.

Sources: S01, S03

Activation & propagation

The electromagnetic coupling obeys the same permission-times-activation rule as other channels. A model of photon propagation additionally needs its kinetic operator. The background activation can affect an interaction coefficient, a kinetic coefficient or both, depending on the specified effective action.

For a transverse mode in a stationary homogeneous readout, a simple quadratic kinetic operator has temporal coefficient A and spatial coefficient B. Positive coefficients give a real characteristic speed. This is a proposed calculation template, not an assigned value for either coefficient.

$$\omega^2 = \frac{B}{A}|\mathbf{k}|^2$$

$$c_{\mathrm{EM}}^2 = \frac{B}{A} \quad (A > 0,\ B > 0)$$

Sources: S01, S03

Why the key matters

The author's argument is that a realised channel imposes the conditions under which an excitation propagates. The photon therefore reads the electromagnetic channel's propagation law. In a regime with stable coefficients, that law supplies a stable characteristic speed.

It is not necessary to assume that every other possible channel has the same kinetic coefficients in every epoch. If the framework predicts equality, the equality should follow from the interaction structure or an effective symmetry. If it predicts a difference, the difference needs its own propagation and detection calculation.

A direct experiment measures arrival times, frequencies, polarisation or interference. The interpretation connects those measurements to A, B and their background dependence.

Sources: S01, S03

Light as a historical messenger

The source argument gives light a further role: it can carry information across changing interaction regimes if the electromagnetic propagation channel remains available along the path. A change in some other sector need not automatically erase every photon.

Survival is a transport question. It depends on the kinetic equation, scattering, absorption, mixing and the history of the background. It is not guaranteed simply by declaring the photon to have one key.

The later photon-history module gives the corresponding transfer calculation. The wider aim is to ask whether observable light can preserve information about interaction configurations different from those inferred from present-day matter.

Continue with: M06 / M35 / M45

Sources: S01, S03

The second one-key example

The gluon is the Strong-key excitation in the schematic account. Its ability to interact with other gluons belongs to that same nonabelian channel.
Self-interaction does not automatically mean that it possesses a second kind of key.

This matters for the central comparison. The photon is not the only one-key example in the author's discussion. Earlier AI passages that say so while listing the gluon as Strong-only are inconsistent. This edition uses the two examples together.

Their shared one-key classification does not make their dynamics identical. The Strong channel's nonabelian structure supplies interactions absent from a free electromagnetic wave.

Sources: S01, S03

One key, nonabelian structure

The effective Yang-Mills field strength contains a derivative part and a commutator part. The latter supplies interactions within the same gauge channel. Permissions identify channel access; the gauge algebra determines the allowed structure inside it.

In a local high-frequency analysis, characteristic propagation is obtained from the principal kinetic operator. Binding, confinement and background interactions affect the physical interpretation and the accessible observables. A freely isolated gluon is not an interchangeable experimental messenger with a photon.

The theory must therefore compare like quantities: a characteristic speed, a group velocity in a specified background, or a measured signal transfer. These need not be inferred by the same procedure.

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + ig[A_\mu, A_\nu]$$

Sources: S01, S03

Binding energy & geometric response

The source discussion distinguishes a direct Higgs key from the contribution of Strong binding to an effective mass or gravitational response. That distinction should remain visible. Energy stored in a field configuration can enter a composite system's stress-energy without assigning the elementary excitation every constituent's permission.

In the author's gravity-as-geometry account, the combined configuration contributes to the emergent geometric readout. The gluon's Strong key then remains its direct channel label. How that contribution is calculated belongs to the effective action and stress-energy map.

This avoids using indirect influence as an unrestricted rule for adding keys. Otherwise every coupled system would rapidly lose the distinctions that KEYCHAIN was introduced to explain.

Sources: S01, S03

The comparison with photons

Photons and gluons provide two questions for the same architecture. Why is each assigned its channel? What fixes the propagation law within that channel? Their comparison tests whether a common speed follows from a shared effective causal structure or whether different regimes can produce distinct characteristic coefficients.

A one-key count is the starting classification. A kinetic calculation supplies the speed. A source, path and detector supply the observation. The distinction allows the author's propagation hypothesis to become a concrete investigation.

The following modules extend the account to multi-key matter, dormant permissions and channel mixing. They keep the photon and gluon examples as anchors throughout the book.

Continue with: M04 / M06 / M37 / M45

Sources: S01, S03

Simultaneous access

An electron is the source's electromagnetic, weak and Higgs permission example. A quark additionally has Strong access. A neutrino has weak access and the mass-related channel specified by the selected neutrino construction.

These examples introduce simultaneous constraints. The effective equations must combine the accessible interactions, rather than replacing one permission by another whenever the particle enters a different environment. Couplings and backgrounds determine which contributions dominate a particular process.

The labels are schematic. Electroweak representations, chirality and mixing still determine the detailed field content. A public keychain diagram organises those questions without being a substitute for their representation matrices.

Sources: S01, S03

Mass coupling & inertia

The simplest source mass template uses a Higgs activation, a state occupation readout and a species coefficient. It is a defined factorisation, not yet a computed table of observed masses. A common background factor can correlate the masses of several species.

For relativistic propagation in a locally constant background with the conventional quadratic dispersion, velocity depends on momentum and the effective mass. The result already allows the speed to change with energy while the permission pattern stays fixed.

This is why adding a key must not automatically be interpreted as imposing an energy-independent fraction of the light speed. The combined kinetic and mass structure determines the relation.

$$m_p = y_p^{(0)} v^{(0)} \alpha_H F_H(P_H)$$

$$v = \frac{c^2 p}{\sqrt{m_p^2 c^4 + p^2 c^2}}$$

Sources: S01, S03

Changes & decay channels

Weak permissions concern allowed transitions as well as propagation. A decay requires an available interaction, an appropriate matrix element and accessible final states. A nonzero permission alone does not calculate a decay rate.

The same structure applies to confinement. Strong access identifies the channel; the nonabelian field dynamics and the physical regime determine binding and accessible excitations. A change in activation could alter those conditions, but the altered spectrum must be solved.

The framework's value is its common language for these changes. The exact quantitative relation between activation and each rate, spectrum or mass remains attached to the selected effective model.

Sources: S01, S03

What can be compared directly

Compare two species sharing a channel while holding their permissions fixed. The channel's activation factor cancels in their effective-coupling ratio. Compare one species at two backgrounds, and the ratio tracks activation if its bare scale and permission are fixed.

These relations are stronger than a general statement that everything changes. They identify combinations that should stay constant and combinations that should drift together. An observation can then challenge a particular factorisation.

Later modules apply this logic to clock comparisons, mass ratios and hidden sectors. It is one of the most direct ways to connect the entire permission architecture with measurements.

Continue with: M12 / M37 / M41 / M43

Sources: S01, S03

Available structure, inactive channel

A dormant key means that the permission exists while the relevant activation is suppressed. It does not mean that the species has been removed from the mathematical state space. The distinction is the basis for the source's hidden-channel and epoch arguments.

The author's propagation proposal asks what remains of a particle's constraints when some channels are inactive. Counting the complete permission list as though every entry were fully engaged would miss that question.

The calculation must retain activation factors wherever the chosen action places them. It can then determine whether dormancy changes interactions, effective inertia, propagation coefficients, detectability or some combination of these.

Sources: S01, S03

What vanishing coupling establishes

In the factorised rule, zero activation removes the corresponding interaction coefficient. That is an exact endpoint statement. It does not specify the remaining kinetic operator or a path through an emergent spacetime.

If the kinetic term survives, an excitation can remain a propagating mode even when an interaction is suppressed. If the proposed spacetime readout itself ceases to apply, a spacetime velocity may cease to be defined. These are different constructions and need different equations.

The zero-coupling endpoint therefore cannot be replaced by division by a zero key count. Such a division is not part of the given coupling rule.

$$\alpha_a = 0 \implies g_{pa} = 0$$

$$\mathbf{v}_p = \nabla_{\mathbf{p}} E_p$$

Sources: S01, S03

Testing the wider claim

The claim that another accessible regime permits propagation faster than the present electromagnetic characteristic speed needs a dispersion relation and a readout between source and detector. It also needs the causal and stability properties of the combined system.

This is a calculational question within the theory, not something decided by a word such as timeless, dormant or higher-dimensional. A vanishing proper-time interval and an infinite signal velocity are not equivalent definitions.

The same standard applies to the source's suggestion of many dormant candidates. The adjoint dimension alone is not a counted catalogue of particles with measured speeds. Their states, kinetic operators and observability have to be assigned.

Sources: S01, S03

Preserve the intended investigation

I retain the author's substantive question: which limits belong to a present active channel, and which belong to the deeper interaction architecture? That question is more general than whether a familiar massive particle approaches a familiar light cone.

The useful next step is to write the coupled kinetic operator, identify its characteristic surfaces and specify how a permitted detector receives a signal. The answer may distinguish propagation from invisibility or from absence of a spacetime trajectory.

This preserves the original direction of inquiry while replacing earlier AI declarations of certain instantaneous propagation with a calculation that can actually decide the case.

Continue with: M06 / M35 / M40 / M45

Sources: S01, S03

The channel-speed hypothesis

The source discussion treats the familiar light speed as a characteristic of the electromagnetic propagation regime. It asks whether a shared deeper structure forces other channels to have the same characteristics, or whether some activation regimes can differ.

This hypothesis should be investigated through coefficients and equations. The current value of a characteristic speed is not derived by assigning the word open to a channel. A comparison between sectors must fix the clock, distance and energy conventions used by the observer.

Photons and gluons are therefore complementary examples. Their one-key classification motivates the comparison; their actual kinetic laws decide it.

Sources: S01, S03

A quadratic propagation template

For real mode amplitudes collected into q, take a homogeneous quadratic action with positive temporal kinetic matrix A and spatial-gradient matrix B. Its principal equation gives a generalised eigenvalue problem. The squared characteristic speeds are the eigenvalues of A inverse B when the required diagonalisation is well defined.

This is an explicitly proposed effective template for the investigation. It introduces no measured coefficients and does not declare a particular channel superluminal. It shows exactly what must be computed from a chosen background.

$$\mathcal{L}_{\mathrm{kin}} = \tfrac{1}{2}\dot{q}^{T} A \dot{q} - \tfrac{1}{2}\nabla q^{T} B \nabla q$$

$$\det(\omega^{2} A - |\mathbf{k}|^{2} B) = 0$$

Sources: S01, S03

Common & different characteristics

If B equals a common squared speed times A, all modes share that characteristic speed. Different interaction strengths can coexist with the same principal cone. If that proportionality fails, the eigenmodes may have different characteristics, subject to stability and the full constraints.

A mass matrix affects finite-frequency dispersion even when the high-frequency characteristics agree. Mixing can also make the propagating eigenmodes differ from the channel-labelled modes. These distinctions are central when comparing one-key and multi-key descriptions.

Thus a permission architecture alone does not require either equal or unequal speeds. It organises the coefficients whose origin the theory must explain.

$$B = c_*^2 A \implies \omega^2 = c_*^2 |\mathbf{k}|^2$$

Sources: S01, S03

A complete propagation experiment

Specify a source excitation, the evolving activation background and the detector's permission operator. Solve the transport equation, then compute the detector readout. Arrival time, frequency and polarisation are outputs of that combined problem.

A locally calculated group velocity is not automatically a signal front or a cosmological separation rate. The calculation must name the quantity being compared. This also keeps cosmic expansion separate from local transport.

The programme links the deepest ontology to a measurable question: how does an interaction state determine the channels through which information can be emitted, propagated and received?

Continue with: M02 / M03 / M35 / M45

Sources: S01, S03

Selecting a subspace

A projector selects a subspace of a vector space. For an orthonormal basis and an index set S_a, the sum of the basis dyads belonging to that set is an orthogonal projector. Acting on a state retains those components and removes the rest.

This is the natural mathematical translation of a sector permission when the sector is a specified internal subspace. A subgroup branching can supply such subspaces, provided the basis and representation action are explicit. The projector's rank is the selected dimension.

A collection of projectors may overlap. Overlap is a statement about their selected subspaces; completeness is a separate statement about their sum. The book does not assume either property merely because the projectors have different sector names.

$$\Pi_a = \sum_{i \in S_a} |e_i\rangle\langle e_i|$$



Selected components

An exact selection, not a particle picture

Sources: S04, S05, S07, S09

The defining identities

An orthogonal projector is Hermitian and idempotent. Hermiticity means that its adjoint equals itself. Idempotence means that applying the selection twice has the same effect as applying it once. Its eigenvalues are zero or one.

For a finite-dimensional orthogonal projector, the trace equals the rank. This identity supports occupation and dimension calculations throughout the manual. It does not determine which subspace corresponds to a physical interaction; that identification remains part of the model construction.

If two projectors are built by selecting subsets of the same orthonormal basis, they commute, even when those subsets overlap. Noncommutativity needs a different relationship between the subspaces and cannot be inferred from overlap alone.

$$\Pi_a^2 = \Pi_a, \qquad \Pi_a^\dagger = \Pi_a$$

$$\mathrm{Tr}\,\Pi_a = d_a$$

Sources: S04, S05, S07, S09

Direct matrix verification

In a basis adapted to the selected sector, the projector has ones on the selected diagonal entries and zeros elsewhere. Squaring the matrix leaves every entry unchanged. Transposing and conjugating it also leaves it unchanged. Counting its diagonal ones gives its trace and rank.

For two common-basis selections, matrix multiplication keeps the intersection of their selected indices. Reversing the multiplication gives the same intersection. Therefore their commutator vanishes. This is a direct finite calculation independent of a physical interpretation.

If their intersection is nonempty, their product is nonzero. The projectors then commute but are not mutually orthogonal. The distinction is relevant when their sum is used as a permission operator.

$$\Pi_a \Pi_b = \Pi_b \Pi_a$$

$$\Pi_a \Pi_b = 0 \quad \text{only for disjoint selections}$$

Sources: S04, S05, S07, S09

Using sector names carefully

The names electromagnetic, weak, strong, Higgs, gravity and dark identify proposed roles. Each role needs the selected subspace and the operator action on the relevant states. A name cannot replace the construction of the projector.

Gauge-generator subspaces and matter-representation subspaces should be distinguished. The Higgs is represented by matter fields with transformation properties; it is not simply another gauge subalgebra. The same care applies to gravity and sterile sectors.

When reviewing a sector definition, ask for its basis, rank, generators, overlap with other sectors and physical readout. Once those are available, the projector can be checked using the identities on these pages.

Continue with: M08 / M13 / M17

Sources: S04, S05, S07, S09

Combining selected sectors

The source formulation combines sector permissions with weights. This is a useful operator construction, but its result is not always an orthogonal projector. The manual writes K_p for the composite so that its algebraic type can be inspected before it is used as a filter or effect.

If the projectors are mutually orthogonal and the weights are only zero or one, their selected sum is again a projector. Fractional weights generally remove idempotence. Overlapping sectors can produce eigenvalues larger than one when they are added without further normalisation.

These are mathematical properties of the construction, not disagreements about the keychain metaphor. They determine which observable formulas are well defined and which extra assumptions are needed.

$$K_p = \sum_a w_{p,a} \Pi_a$$

Sources: S04, S05, S07, S09

Squaring the composite

Expand the square rather than assuming the projector identity. The expansion contains weighted diagonal terms and cross terms. With orthogonal projectors, the cross terms vanish, but the squared weights remain. Idempotence then requires each nonzero weight to equal one.

With overlapping projectors, cross terms contribute. If all selected projectors are positive and weights are nonnegative, the sum is positive. Positivity alone does not make it bounded above by the identity. A measurement effect requires that additional bound.

An internal permission operator may be useful without being a measurement effect. The book therefore names the intended use before choosing an expectation or normalisation formula.

$$K_p^2 = \sum_{a,\,b} w_{p,\,a} w_{p,\,b} \Pi_a \Pi_b$$

$$0 \le E_p \le I \quad \text{for a measurement effect}$$

Sources: S04, S05, S07, S09

An exact fractional-weight check

Take a nonzero projector Pi and multiply it by one half as an operator test. The square is one quarter times Pi, so it is not equal to the original operator. Its eigenvalues are zero and one half. It is a positive effect, but it is not an idempotent projector.

This finite example tests the source allowance of fractional weights. It assigns no measured physical coupling and introduces no fitted model constant. Its purpose is to show why a weighted species permission needs its own algebraic label.

If two identical permissions are added with unit weights, the result is twice Pi. It remains positive, but its selected eigenvalue is two and it cannot be a probability effect without another definition.

$$K = \tfrac{1}{2}\Pi \quad \Longrightarrow \quad K^2 = \tfrac{1}{4}\Pi$$

Sources: S04, S05, S07, S09

Consequences for observables

For a filter K acting on a state, the norm of the filtered state uses K dagger K. For a measurement effect E, a compatible elementary filtering operator is its positive square root, subject to the chosen measurement instrument. These constructions use different denominators.

The next modules retain this distinction. A composite keychain can remain a permission label while the readout uses a separately defined effect. This prevents the same weighted operator from being treated as three different objects without explanation.

When adapting an older equation, change the operator type and normalisation together. A notation change alone cannot fix the meaning of an expectation value.

Continue with: M09 / M11 / M17

Sources: S04, S05, S07, S09

A probability obtained from a state

Occupation of a sector is calculated from a normalised state and an orthogonal projector. The result is real, nonnegative and no greater than one. It answers how much of the specified state lies in the selected subspace.

Occupation is not the same as having permission to interact. A species permission is a model assignment; occupation depends on the state. Nor is occupation automatically an independent local activation field. The book writes P_a for occupation to keep these roles separate.

For a mixed state, replace the vector expectation by the trace of the density matrix with the projector. This permits an ensemble description while retaining the same probability interpretation.

$$P_a = \langle \Psi | \Pi_a | \Psi \rangle$$

$$P_a = \mathrm{Tr}(\rho \Pi_a)$$

Sources: S04, S05, S07, S09

Completeness changes the constraints

If the projectors are mutually orthogonal and sum to the identity, their occupations sum to one. This follows directly from state normalisation. The individual occupations are therefore constrained and cannot all vanish in the same normalised state.

If projectors overlap, the occupations need not form a mutually exclusive probability distribution. If the collection is incomplete, their sum need not be one. The probability of each orthogonal projection remains meaningful, but the interpretation of the whole list changes.

These conditions must be written before using sector occupations as cosmological switches. Independent activation fields can all be assigned zero in a model, whereas a complete set of mutually exclusive occupations cannot.

$$\sum_a \Pi_a = I \quad \Longrightarrow \quad \sum_a P_a = 1$$

Sources: S04, S05, S07, S09

The uniform-state calculation

The source defines a uniform superposition over an orthonormal basis. For a sector projector selecting d_a of those basis directions, the expectation is d_a divided by 248. Each selected component contributes the same squared amplitude.

This result depends on the projector being diagonal in the basis in which that particular uniform state was defined. A uniform list of coefficients in one basis need not have equal amplitudes in another basis. The state and the basis convention must travel together.

The completely mixed state I divided by 248 gives the same rank-based occupation for any rank-d_a projector. That basis-independent ensemble is different from a chosen pure uniform superposition.

$$|\Psi_0\rangle = \frac{1}{\sqrt{248}} \sum_i |e_i\rangle$$

$$\rho_* = \frac{I}{248}, \qquad P_a = \frac{d_a}{248}$$

Sources: S04, S05, S07, S09

Keeping occupation and activation separate

If a model sets alpha_a equal to P_a, it inherits the occupation constraints and state dependence. If alpha_a is an independent scalar field, it needs an independent action or evolution law. A hybrid formulation must specify the relationship rather than alternating between the definitions.

The distinction affects claims about all sectors becoming inactive, the response of a Hamiltonian and the proposed mass sum. It also affects linearity: placing a state-dependent expectation inside H can make the effective Schrödinger equation nonlinear.

Use this module to identify which activation version a later calculation assumes. The activation section records both constructions and their consequences.

Continue with: M10 / M19 / M20

Sources: S04, S05, S07, S09

M10 / 1 OF 4

Permission and local availability

V.A.S.K. makes channel availability part of the changing physical configuration. In this effective-field formulation alpha is a local availability field. The alternative occupation-derived formulation is retained explicitly in M19.

Activation supplies the proposed local modulation of a permitted interaction. In the independent-field formulation, alpha_a assigns a real number between zero and one to each effective spacetime event. Its value can change while the species permission remains fixed.

This formulation expresses the KEYCHAIN distinction between carrying a key and finding an active lock. The model's dynamics must then determine alpha rather than leaving it as an arbitrary slider. The action, potential, couplings and conditions specify that task.

An occupation-defined activation is an alternative construction. It derives its value from the state and inherits the constraints described in the preceding module. The manual does not merge the two versions into one unstated definition.

Sources: S04, S05, S07, S09

The independent-field definition

Let the effective spacetime base be M and define alpha_a as a map from M to the closed unit interval. The coupling product then uses this field at the same event as the effective matter and gauge variables. A spacetime-dependent coupling requires attention to gauge invariance in the full action.

A local field has a kinetic structure and interacts with other fields through specified terms. These terms determine whether constant activation is stable, whether gradients cost energy and whether transitions can occur. Assigning a range does not establish those properties.

The unit interval can be enforced through a constrained theory or a parametrisation. The choice affects field equations, endpoint behaviour and numerical evolution, so it belongs in the model specification.

$$\alpha_a : \mathcal{M} \longrightarrow [0, 1]$$

$$g_{pa}(x) = g_a^{(0)} w_{p,a} \alpha_a(x)$$

Sources: S04, S05, S07, S09

Endpoint and ratio checks

The product rule gives a zero effective coupling at zero activation and the permitted bare coupling at unit activation. This is a direct limiting check. It does not show which endpoint is reached or whether a potential makes it stable.

A ratio of activations at two epochs can exceed one even though both activations lie in the unit interval. The numerator only needs to exceed the denominator. Therefore a ratio claim must not be rewritten as an absolute activation outside the allowed range.

If a mass is assumed proportional to activation, the same caution applies to mass ratios. The full mass formula may include a changing vacuum scale or other factors, so the proportionality and its regime must be stated.

$$\frac{g_{pa}(x_1)}{g_{pa}(x_2)} = \frac{\alpha_a(x_1)}{\alpha_a(x_2)}$$

Sources: S04, S05, S07, S09

The next dynamical question

Once the activation definition is chosen, ask what its equation predicts. A potential identifies stationary configurations. A kinetic term sets the response to gradients. Initial conditions determine which solution is followed. Couplings to other fields provide back-reaction.

The proposed cosmological epochs correspond to activation remaining near selected configurations for long periods. To calculate their duration or sequence, the dynamics and transition rates need to be evaluated. A drawing of successive plateaus is a useful guide to this calculation, not its output.

The activation modules now develop these obligations using the supplied potential variants and exact derivatives. Their status remains tied to the stated inputs.

Continue with: M19 / M22 / M33

Sources: S04, S05, S07, S09

Conditioning a state

A sector-conditioned observable asks for an expectation after a specified selection or filtering operation. For an orthogonal projector, the filtered state is Pi times Psi. Its squared norm gives the probability of the selected sector. Dividing by that norm produces the conditional expectation when the probability is nonzero.

For a general filtering operator K, the squared norm contains K dagger K. The denominator must match the operation used in the numerator. Replacing it with K alone can give an incorrect result even for the identity observable.

A measurement effect E specifies an outcome probability, but a full measurement instrument also specifies the state update. Choosing the positive square root is one elementary construction, not the only possible instrument.

$$\langle O \rangle_K = \frac{\langle \Psi | K^\dagger O K | \Psi \rangle}{\langle \Psi | K^\dagger K | \Psi \rangle}$$

Sources: S04, S05, S07, S09

Projector and effect versions

When K is an orthogonal projector, K dagger K equals K. The familiar projector denominator is then correct. This special case explains why the formula works in the simplest sector construction but cannot be transferred unchanged to every weighted keychain.

For a positive effect E bounded by the identity, the outcome probability is the expectation of E. If the filtering operator is its positive square root, the numerator contains square roots of E on both sides of the observable. The denominator is the expectation of E.

The formula must state which selection is used. A reader should never have to guess whether a species operator is an idempotent projector, a general filter or a measurement effect.

$$K = \sqrt{E} \quad \Longrightarrow \quad K^{\dagger} K = E$$

Sources: S04, S05, S07, S09

The identity-observable check

Set O equal to the identity. A correctly normalised conditional expectation must be one whenever the denominator is nonzero. In the general filter formula, the numerator and denominator are identical, so the result is one exactly.

This provides a fast consistency test for any proposed conditioning rule. With a fractional permission substituted into the projector-only denominator, the result can differ from one. That failure identifies a normalisation mismatch, not a novel physical identity observable.

If the selection probability is zero, the conditional state is undefined. A zero denominator cannot be treated as evidence for an infinite expectation or instantaneous behaviour. The selected event has no normalised conditional state under that rule.

$$O = I \quad \Longrightarrow \quad \langle I \rangle_K = 1$$

Sources: S04, S05, S07, S09

Using a readout in the model

The readout rule connects internal operators to predictions. It must therefore be treated as carefully as the Hamiltonian. State the operator, the state, the selection and the normalisation. If the observable is position-dependent, also state the effective field description supplying that dependence.

Different filters can produce different conditioned states while sharing an outcome probability. This is why a probability effect alone does not determine all subsequent dynamics. A complete measurement construction needs the instrument or another specified state-update rule.

The source's particle permissions can remain useful organising objects. The public manual improves their calculational use by separating permission, filtering and observation wherever their mathematical properties differ.

Continue with: M08 / M09 / M18

Sources: S04, S05, S07, S09

A consequence of factorisation

The coupling rule does more than suppress a channel at zero activation. Under fixed bare scales and fixed species permissions, every species sharing sector a receives the same multiplicative activation factor. This gives an immediate set of correlated changes and invariant ratios.

Take two species with nonzero permission in the same sector. Their effective coupling ratio is independent of activation. Compare one species at different locations or times, and the coupling ratio follows the activation ratio if the other inputs remain fixed.

These are conditional results of the stated rule. They are suitable targets because a future calculation or observation can test the factorisation itself.

Sources: S04, S05, S07, S09

Ratios & drift

Differentiate a nonzero coupling while holding its bare scale and permission fixed. Its logarithmic derivative equals the logarithmic derivative of activation. The same equality holds for any other species sharing the sector under those assumptions.

If a dimensionless fine-structure parameter is proportional to the square of that coupling in a fixed convention, its logarithmic drift is twice the coupling drift. This uses a specified low-energy identification; it is not a replacement for renormalisation.

$$\frac{g_{pa}}{g_{qa}} = \frac{w_{p,a}}{w_{q,a}}$$

$$\partial_\mu \ln g_{pa} = \partial_\mu \ln \alpha_a$$

$$\partial_\mu \ln \alpha_{\mathrm{fs}} = 2\partial_\mu \ln \alpha_{\mathrm{EM}}$$

Sources: S04, S05, S07, S09

What a deviation would mean

A varying same-sector species ratio would require a change in the permissions, bare scales, representation assignment or the factorised coupling rule. The framework therefore has a concrete place to locate a discrepancy. It need not absorb every outcome by relabelling activation.

Ordinary energy-scale running must be treated separately from drift across a changing background. Compare measurements at a common scale or evolve them with the stated renormalisation prescription. Otherwise different causes of variation become numerically entangled.

The equations remain useful when additional structure is introduced: the new structure must state which assumption it changes and predict the resulting departure.

Sources: S04, S05, S07, S09

Mass ratios & identifiable inputs

The simplest mass template also has a common background factor. If two species share the same Higgs activation and the same occupation-dependent factor, their mass ratio equals the ratio of their species coefficients. A changing common Higgs background then changes both masses together.

This identifies an important limit of the template. It does not independently explain a changing mass ratio unless another species-dependent contribution is present. It also does not derive the coefficients merely by writing them as overlap symbols.

A predictive calculation must separate what is fixed by measured input from what is obtained independently. The resulting correlations can then be used across propagation, clocks and cosmology.

$$\frac{m_p}{m_q} = \frac{y_p^{(0)}}{y_q^{(0)}}$$

Continue with: M04 / M19 / M41 / M43

Sources: S04, S05, S07, S09


The internal substrate

The selected internal substrate is the complexified adjoint representation of E8. Its dimension is 248. Eight commuting Cartan directions organise the root-space description, and the remaining generator directions correspond to 240 roots. This is a statement about Lie-algebra structure.

The model uses this structure to organise sector permissions. A root is a vector in an eight-dimensional Euclidean description; a generator is an element of the algebra; a state is a vector in the chosen representation space. These objects are related, but they are not identical.

Naming each object prevents a diagram of roots from being mistaken for a physical arrangement of particles. The physical interpretation requires representation assignments, effective fields and a map to observed interactions.

$$\dim \mathfrak{e}_8 = 240 + 8 = 248$$



First two root coordinates; coincident images merged

Sources: S04, S05, S07, S09

An explicit root construction

Use eight orthonormal Euclidean coordinate directions. One root family has two entries equal to plus or minus one and the other six entries zero. Choosing the two positions gives 28 choices, and their four sign choices give 112 roots.

The second family has all eight entries equal to plus or minus one half, with an even number of minus signs. There are 128 allowed sign patterns. Every vector in either family has squared length two. Together they supply the 240 roots.

This construction is suitable for a finite code check. Keep exact integer coordinates by doubling the vectors, then divide dot products by four when returning to the conventional normalisation.

$$4\binom{8}{2} = 112, \qquad 2^7 = 128$$

$$112 + 128 = 240$$

Sources: S04, S05, S07, S09

Root lengths and angles

For the first family, the two nonzero unit entries give squared length two. For the second family, eight squared halves sum to two. The common length makes the angle between roots depend directly on their dot product.

For distinct non-opposite roots in this normalisation, the possible dot products are minus one, zero and one. These give angles of 120, 90 and 60 degrees. Opposite roots have angle 180 degrees. This finite set concerns roots, not all possible vectors of the lattice or their spatial projections.

A very small continuous misalignment used elsewhere in the model therefore needs another construction. It cannot be described as the smallest angle between distinct E8 roots without changing the object under discussion.

$$\cos\theta = \frac{r \cdot s}{2}$$

Sources: S04, S05, S07, S09

From a root list to an algebra

The root list supplies geometry and representation weights. A Lie algebra also needs its bracket and the normalisations of its generators. Assigning a linear transformation to roots is not enough to establish that it preserves all brackets.

An algebra automorphism must preserve the bracket for every pair of elements. A candidate Z action can therefore be checked at two levels: its finite linear action on the representation, and its compatibility with the algebraic multiplication. These are distinct calculations.

The Z section uses the root and branching structure as an organised substrate. It keeps the automorphism-identification obligation explicit where a physical or algebraic interpretation requires it.

Continue with: M14 / M15 / M48

Sources: S04, S05, S07, S09

Subgroups reorganise the same space

A branching rule describes how a representation decomposes when restricted to a subgroup. The total dimension is preserved. The resulting blocks may transform differently under that subgroup, even though they originated within one irreducible representation of the larger group.

This is the basis for defining candidate sectors in V.A.S.K. The projectors select representation blocks after a branching has been specified. Different branching routes organise the space differently; they require a correspondence before their labels can be combined.

Quadratic Casimirs depend on the algebra, representation and normalisation. A value for the full E8 adjoint must not silently be replaced by a value for an SO10 vector or a restricted subgroup operator.

$$C_2(R) I_R = \sum_A T_R^A T_R^A$$

Algebra | Adjoint dimension | C2 adjoint

U1 | 1 | 0: ratio undefined

SU2 | 3 | 2

SU3 | 8 | 3

G2 | 14 | 4

F4 | 52 | 9

E6 | 78 | 12

E7 | 133 | 18

E8 | 248 | 30

Sources: S04, S05, S07, S09

Representation and adjoint values

The source uses the conventional adjoint values two for SU2, three for SU3 and thirty for E8. A matter representation has its own value: the SU2 fundamental has three quarters and the SU3 fundamental has four thirds in the corresponding conventions.

For SO10 with vector-index normalisation one, the vector 10 has Casimir nine halves, and the chiral spinor 16 has Casimir forty-five eighths. Their dimension-to-Casimir ratios are therefore different from ratios built using an adjoint Casimir.

The U1 adjoint is neutral and its quadratic adjoint Casimir is zero. Dividing its dimension by that zero does not define a finite stability ratio. Any special U1 baseline is an additional model convention.

$$C_2(10) = \frac{9}{2}, \qquad C_2(16) = \frac{45}{8}$$

Sources: S04, S05, S07, S09

An exact ratio comparison

For the stated SO10 representation convention, the vector ratio is ten divided by nine halves. The spinor ratio is sixteen divided by forty-five eighths. Performing the arithmetic gives twenty ninths and one hundred twenty-eight forty-fifths.

The spinor ratio exceeds the vector ratio by twenty-eight forty-fifths. Consequently a rule favouring the smallest dimension-to-Casimir ratio does not select the spinor ahead of the vector under this particular definition. The result checks the definition; it does not choose a physical activation model.

If another restricted operator is intended, state that operator and repeat the calculation. A change of definition must not be presented as an unchanged numerical consequence of the old formula.

$$\mathcal{R}_{10} = \frac{20}{9}, \qquad \mathcal{R}_{16} = \frac{128}{45}$$

$$\mathcal{R}_{16} - \mathcal{R}_{10} = \frac{28}{45}$$

Sources: S04, S05, S07, S09

Inspecting a Casimir filter

Before using a Casimir in an activation inequality, identify which generator sum appears and on which representation it acts. State the dimension counted by d_a. A subalgebra dimension and the dimension of a matter block answer different questions.

Then fix the threshold's temperature and coupling normalisation. Ratios can order candidate thresholds, but they do not fix which sectors are active at a particular epoch without the actual cutoff and dynamics. Conjugate representations also share the same quadratic Casimir.

The later threshold and dormancy modules apply these checks to the source formulas. This preserves the filter as a proposed organising mechanism while making its current calculation reproducible.

Continue with: M16 / M24 / M42

Sources: S04, S05, S07, S09


A selected branching route

The proposed generation construction uses the E6 x SU3 branching of the E8 adjoint. The decomposition contains an E6 adjoint, an SU3 adjoint, a tensor block transforming as 27 times 3, and its conjugate. The dimensions add to 248.

The SU3 factor in this route is labelled generation SU3 to distinguish it from colour SU3. The model chooses a cyclic element acting on the three-component generation index. Its group membership does not decide whether the symmetry is gauged, global or explicitly broken in the effective dynamics.

The branching provides a candidate location for three selected SO10 spinor copies. It also contains other representations and conjugate blocks whose physical roles require a full spectrum assignment.

$$248 = 78 + 8 + 81 + 81$$



Sources: S04, S05, S07, S09

The selected 16-dimensional copies

Under Spin10 x U1, the E6 fundamental 27 decomposes into a chiral spinor 16, a vector 10 and a singlet. The conventional charges in the supplied route are one, minus two and four, respectively.

Tensoring the spinor component with the generation triplet supplies three copies of the internal 16. Selecting those copies defines the 48-dimensional generation block used in the explicit Z construction. It does not remove the other components of the 27 or the conjugate 81-dimensional block.

Spin10 is used here for the group that carries spinor representations. Its Lie algebra is the same as that conventionally labelled so10. The distinction avoids suggesting that the internal spinor is an ordinary vector representation.

$$27 = 16_1 \oplus 10_{-2} \oplus 1_4$$

$$\mathcal{H}_{\text{gen}} = \mathbb{C}^3 \otimes \mathbb{C}^{16}$$

Sources: S04, S05, S07, S09

Dimension and action checks

The selected block has dimension three times sixteen. The full 27 times 3 block has dimension eighty-one. Its thirty-three remaining directions belong to the triplets of the vector 10 and singlet 1. They are part of the full generation-SU3 action.

The E6 adjoint block is fixed by a subgroup element acting only on generation SU3. The SU3 adjoint transforms by conjugation, and the conjugate triplet transforms with the conjugate action. These details determine the full 248-dimensional spectrum later in the book.

The dimension sum is a necessary consistency check. Identifying the selected copies as the observed chiral families additionally requires Lorentz assignments, gauge charges and a demonstrated treatment of the other states.

$$81 - 48 = 33$$

Sources: S04, S05, S07, S09

Physical questions left by the branching

The branching does not itself choose an active vacuum, derive Yukawa matrices or eliminate mirror degrees of freedom. A permission or activation mechanism may be proposed for these tasks, but it must specify the operators and the resulting effective spectrum.

The model states that generation SU3 is global in its intended construction. An explicitly chosen Hamiltonian can then break that symmetry. If a gauged version is later introduced, gauge invariance, gauge fields and anomaly conditions would need to be reconsidered.

This route gives the Z section a precise internal space and a candidate full action. Keeping those facts separate from the physical completion makes the construction both readable and reviewable.

Continue with: M26 / M28

Sources: S04, S05, S07, S09

A second organisation of the substrate

The source also uses a route through so16 to so10 plus so6. At the group level, spin covers are required when spinor representations are used. The E8 adjoint decomposes into the so16 adjoint and one chiral spinor of Spin16.

The so16 adjoint has dimension 120, and the selected spinor has dimension 128. Under Spin10 x Spin6, the adjoint contains blocks of dimensions 45, 15 and 60. The spinor supplies two tensor blocks of dimension 64 each.

This route is useful for inspecting the source's unwanted-sector discussion. It must be related explicitly to the E6 x SU3 route before labels from both decompositions are treated as one common sector list.

$$248 = 120 + 128$$

$$120 = 45 + 15 + 60$$

Sources: S04, S05, S07, S09

The representation blocks

The adjoint branches as (45,1), (1,15) and (10,6). The spinor branches as a 16 paired with a Spin6 spinor 4 and the conjugate pair. The sum of all five block dimensions remains 248.

These are representation blocks. Calling a block a mirror-fermion sector additionally requires a Lorentz-spin and chirality assignment. The vector 10 and singlet labels in another branching are also not, by their names alone, opposite-chirality copies of the observed families.

The internal charges can be calculated from the chosen subgroup action. Their interpretation as particles needs an effective field construction. The manual keeps that extra step visible when considering dormancy.

$$248 = 45 + 15 + 60 + 64 + 64$$

Sources: S04, S05, S07, S09

The full-adjoint trace check

The supplied dormancy appendix defines an effective Casimir by tracing the full E8 Casimir over a projected block and dividing by its dimension. In the stated adjoint normalisation, the full operator equals thirty times the identity.

Consequently the trace over a rank-d projector is thirty times d. Dividing gives thirty for every selected block. The (10,6) block therefore yields ratio sixty over thirty, and the (1,15) block yields fifteen over thirty.

This calculation supplies the exact result of the written formula without a computer algebra system. It does not establish dormancy. A restricted-subgroup Casimir is a different operator and needs a separately stated definition.

$$\frac{\mathrm{Tr}(\Pi_a 30 I)}{d_a} = 30$$

$$\mathcal{R}_{(10,6)} = 2, \qquad \mathcal{R}_{(1,15)} = \tfrac{1}{2}$$

Sources: S04, S05, S07, S09

Diagonal colour needs an actual restriction

If physical colour is proposed as a diagonal subgroup of two SU3 factors, restrict each representation to that subgroup explicitly. A representation charged under only one factor does not become a singlet merely because the subgroup is diagonal.

For example, an element g acts on a tensor representation R times 1 as R(g) times 1. The same R action survives under the diagonal embedding. A separate projection could remove states, but that projection would need to be defined and justified.

The unwanted-spectrum problem therefore needs both the representation restriction and the activation or removal mechanism. A dimension list or a placeholder ratio cannot replace these calculations.

$$(R,1)|_{\mathrm{diag}} \simeq R$$

Continue with: M14 / M24 / M42

Sources: S04, S05, S07, S09

From a stated sector to a matrix

Let B contain basis vectors for a selected subspace as its columns. If those columns are linearly independent, their Gram matrix B dagger B is invertible. The corresponding orthogonal projector can then be constructed without assuming that the columns were already orthonormal.

This formula is useful for turning a branching or weight-space selection into a finite matrix. It makes the sector definition inspectable: the subspace is carried by B, while the inverse Gram matrix supplies the required normalisation.

If the columns are dependent, the ordinary inverse does not exist. A rank-revealing method or a correctly defined pseudoinverse is required. A numerical routine must report that distinction rather than silently ignoring it.

$$\Pi = B(B^{\dagger}B)^{-1}B^{\dagger}$$

Sources: S04, S05, S07, S09

The defining properties

The inverse Gram matrix is Hermitian when B dagger B is positive definite. Taking the adjoint of the projector expression therefore leaves it unchanged. Multiplying the expression by itself cancels the central Gram matrix against its inverse and returns the original expression.

The projector acts as the identity on the column space of B. It removes vectors orthogonal to that space. Its rank equals the number of independent columns. These facts connect the matrix to the intended selected sector.

The result depends only on the subspace, not on a particular independent basis for it. An invertible change of basis among the columns produces the same orthogonal projector.

$$\Pi^{\dagger} = \Pi, \qquad \Pi^{2} = \Pi$$

Sources: S04, S05, S07, S09

A symbolic verification

Write G for B dagger B. In the square of the projector, the middle factors B dagger B form G. The product G inverse times G times G inverse simplifies to G inverse. The remaining outer factors give the original projector.

For an orthonormal column set, G is the identity. The formula reduces to B B dagger, or the sum of the selected basis dyads. This matches the common-basis sector expression used earlier in the book.

A numerical check should compute the Hermiticity residual, idempotence residual and trace. Residuals are assessed against an explicitly chosen precision and matrix norm. A small residual verifies the implementation of the selected projector, not its physical assignment.

$$\Pi^2 = BG^{-1}GG^{-1}B^\dagger = \Pi$$

Sources: S04, S05, S07, S09

A reproducible construction record

Record the basis columns, their ordering, the generator normalisation and the method used to determine rank. Store the matrix or enough data to regenerate it. List any tolerance used by a numerical decomposition.

If two sector projectors are compared, calculate their product and commutator. This reports overlap and compatibility directly. Do not infer either from the sector names. If a complete decomposition is intended, also verify that the projectors sum to the identity.

This record is the practical basis of an IETM-style inspection module. It states the input, operation, expected identities and interpretation of the output so another reader can repeat the construction.

Continue with: M07 / M08 / M42

Sources: S04, S05, S07, S09

A state requires a dynamical context

A vector in the internal block is a proposed internal state. Its components depend on the chosen basis. A density matrix describes a pure state or an ensemble. Both descriptions require an evolution law and observables before they make predictions.

The finite internal block is not a full Fock space of a quantum field theory. Field excitations over spacetime involve additional degrees of freedom and a statistics assignment. A derivation of the effective field theory must construct those structures rather than infer them from the number 248.

This module sets the internal-state conventions used by the Hamiltonian and Z-gate calculations. It leaves the physical field-theory completion to the later matter and testing modules.

$$\langle \Psi | \Psi \rangle = 1$$

$$\rho \geq 0, \qquad \mathrm{Tr}\, \rho = 1$$

Sources: S04, S05, S07, S09

Evolution with a Hermitian generator

For a state-independent Hermitian Hamiltonian, the Schrödinger equation generates norm-preserving evolution. A time-independent generator gives a unitary exponential. A time-dependent generator requires time ordering when its values at different times do not commute.

The source uses proper time in an effective geometry. A covariant derivative may include a connection. Its compatibility with the state inner product must be stated so that the evolution preserves the intended norm. A connection coefficient does not automatically have the energy units of a Hamiltonian term.

The generation Hamiltonian is one part of the supplied total generator. Setting that part to zero does not imply that the locking, mixing, geometric or bath terms vanish.

$$i\hbar \frac{d|\Psi\rangle}{d\tau} = H|\Psi\rangle$$

$$U(\tau) = e^{-iH\tau/\hbar}$$

Sources: S04, S05, S07, S09

The density-matrix check

For the same closed, linear unitary dynamics, the density matrix evolves by a commutator with H. Taking its trace gives zero because the trace of a commutator vanishes. Thus the state remains normalised.

Unitary evolution also preserves the eigenvalues of the density matrix in a closed system. A reduced state obtained by tracing over a bath can behave differently because its dynamics includes correlations and exchange with that bath. The enlarged description must specify the bath and its coupling.

An expectation inserted into H can make the effective equation state-dependent. Even if the norm is preserved, further requirements such as linearity and causal consistency need separate analysis.

$$i\hbar\dot{\rho} = [H, \rho]$$

$$\mathrm{Tr}[H, \rho] = 0$$

Sources: S04, S05, S07, S09

Choosing a readout

Specify the observable operator and the state before evaluating an expectation. If the calculation is conditional, also specify the filter or effect and its normalisation. If it concerns a physical detector, provide the map from the internal observable to the detector quantity.

The source's sector expectations are finite and calculable once the inputs are given. Predicting an actual cross-section, transition rate or mass requires the effective interaction and kinematic structure. The internal expectation is one ingredient in that larger calculation.

This layered description is useful precisely because each layer can be checked. It becomes a physical prediction when the maps between them are explicit and their approximations are controlled.

Continue with: M11 / M20 / M37

Sources: S04, S05, S07, S09

Choosing the activation construction

The activation field represents the model's local availability of an interaction. Its range is the unit interval. The independent-field construction gives it a separate spacetime dynamics, while the occupation construction derives it from the internal state. A calculation must choose which construction it uses.

Independent fields can all be inactive without violating the normalisation of an unrelated internal state. Complete orthogonal occupations cannot all vanish. This difference matters for the proposed dormant-sector and black-hole interpretations.

This edition uses independent alpha fields when analysing the supplied scalar actions. It uses P_a when evaluating sector occupation from a normalised state. A relationship between the two is an additional model assumption that must be written before it is used.

$$0 \leq \alpha_a(x) \leq 1$$

Sources: S04, S05, S07, S09

Enforcing the allowed interval

An evolution law must keep activation in its allowed domain. One approach uses constrained variables with boundary conditions. Another writes activation as a bounded function of an unconstrained field. The second approach changes the effective kinetic term through the derivative of that function.

A potential with a logarithmic entropy term has endpoint behaviour different from a simple polynomial. Numerical evolution must handle the domain explicitly rather than stepping across a logarithm's singular boundary. A finite time step is not a physical transition rule.

The chosen domain treatment affects stationary points, fluctuations and the meaning of a fully active endpoint. Record it alongside the action so that analytical and numerical calculations use the same model.

CHECK: State how the field equation preserves the activation domain.

Sources: S04, S05, S07, S09

Occupation-defined activation

If alpha_a is set equal to the expectation of Pi_a, it lies in the unit interval for an orthogonal projector and a normalised state. Its evolution follows from that state's dynamics. For a complete orthogonal decomposition, the alpha values then sum to one.

Placing those expectations inside H makes the effective generator state-dependent. The resulting evolution must be analysed as such. Norm preservation can still hold for a Hermitian state-dependent operator, but that fact alone does not establish a fundamental linear theory.

The source proposes an auxiliary bath to realise effective feedback in an enlarged linear system. That proposal needs the enlarged Hamiltonian and reduction rule, which are treated in the energy-balance module.

$$\alpha_a = P_a \quad \Longrightarrow \quad \sum_a \alpha_a = 1$$

Sources: S04, S05, S07, S09

Reading an epoch statement

An epoch is proposed as a regime in which relevant activations remain near a configuration for an extended duration. Constant activation is a description of that regime; it is not a calculation of its stability. The potential and kinetic terms must supply the latter.

The labels Higgs ON and Higgs OFF can identify intended regimes. They do not determine the background scale factor, transition history or survival of earlier photons. A cosmological solution is needed before those observational claims are evaluated.

This module gives the fields a clear domain and interpretation. The remaining activation modules now examine dynamics, stationary points and the threshold claims rather than prescribing an epoch sequence by label.

Continue with: M09 / M22 / M33

Sources: S04, S05, S07, S09

Locking & transition are different

The source locking Hamiltonian assigns energetic preferences to sector occupations. A mixing operator connects selected subspaces. The distinction matters: a term that changes relative phases does not necessarily transfer population.

For fixed projectors, occupation changes through a commutator with the effective generator. If the locking term is made only from mutually commuting sector projectors, it commutes with every one of those projectors. On its own it therefore leaves their occupations unchanged.

This does not remove locking from the framework. It identifies the additional dynamics required for relaxation towards an energetic preference: noncommuting terms, coupling to an auxiliary system, or a separately evolving effective activation field.

Sources: S04, S05, S07, S09

The occupation equation

For unitary state evolution and a fixed projector, differentiating its expectation gives the commutator equation. If the projector moves with time, its explicit derivative contributes too. A covariant description must include the corresponding transported-basis terms.

The result supplies a direct inspection of any claimed activation mechanism. Build the selected operator and evaluate the commutator. A nonzero threshold label is not a substitute for a nonzero matrix element connecting the relevant subspaces.

$$\dot{P}_a = \tfrac{i}{\hbar}\langle [H, \Pi_a] \rangle + \langle \dot{\Pi}_a \rangle$$

$$[H_0, \Pi_a] = 0,\ \dot{\Pi}_a = 0 \implies \dot{P}_a = 0$$

Sources: S04, S05, S07, S09

Exactly dormant & nearly dormant

In the closure alpha equals occupation, mixing multiplied by both sector occupations can leave an exactly empty sector invariant. If every coupling into that sector contains its zero activation, that restricted evolution has no term that can populate it.

A nearly dormant state is different: a nonzero seed can permit subsequent mixing. An ungated auxiliary coupling can also change the endpoint behaviour. Either construction must be included explicitly in the initial state or generator.

In the independent-field closure, the field equation determines whether a zero activation begins to move. Its potential slope and cross-couplings can supply a driving term even when the microscopic occupation is zero.

Sources: S04, S05, S07, S09

A route to persistent epochs

Closed unitary dynamics preserves purity and is reversible. Persistent effective ordering and a directional relaxation history require a stated reduced description, coarse-graining, boundary condition or auxiliary-system mechanism.

The R5 sources provide a place for that mechanism through the enlarged system. The next calculation is to specify the system-bath interaction and derive the reduced equation rather than assuming that every Hermitian locking term produces friction.

This result connects the entire architecture: KEYCHAIN selects access; V.A.S.K. supplies availability; the generator or field action must explain how availability actually changes and stabilises.

Continue with: M19 / M21 / M24 / M40

Sources: S04, S05, S07, S09

The supplied generator structure

The source splits the internal generator into locking, sector mixing, geometry or connection, bath completion and generation mixing. Each term requires a defined operator and units. The sum is meaningful only when all terms act on the specified space and share energy units.

A locking term assigns energies to selected sectors. A mixing term transfers amplitude through off-diagonal operators. A connection describes parallel transport in a chosen effective geometry. A bath term acts on an enlarged space. The Z term acts on the selected generation index.

These terms provide a model architecture. Their coefficients and physical mapping remain essential inputs. Listing the terms does not determine their numerical strengths or prove that they recover a complete effective field theory.

$$H = H_0 + H_{\mathrm{mix}} + H_{\mathrm{gen}} + H_{\mathrm{bath}}$$

Sources: S04, S05, S07, S09

Covariant derivative bookkeeping

The source uses a derivative containing a connection Gamma_tau. Written as d/dtau plus Gamma_tau, it has inverse-time units. Moving that connection to the right-hand side of the Schrödinger equation produces a term with an explicit factor of minus i hbar.

Thus a connection matrix is not automatically a Hamiltonian in energy units. If it is represented as a geometric Hamiltonian contribution, show the conversion. The inner-product compatibility condition determines the appropriate adjoint relation in the representation being used.

For a Lorentz spinor description, the spacetime spinor inner product and frame structure require their own treatment. Internal Euclidean unitarity should not silently be substituted for that covariant construction.

$$D_\tau = \partial_\tau + \Gamma_\tau$$

$$i\hbar \partial_\tau \Psi = (H - i\hbar \Gamma_\tau)\Psi$$

Sources: S04, S05, S07, S09

Mixing operators and units

A Hermitian sector-mixing term can be formed from an operator M_ab plus its adjoint, multiplied by a real coefficient with suitable energy units. If the coefficient is dimensionless, the operator or an additional scale must supply the energy dimension.

The source's temperature and gravitational-potential factors are dimensionless ratios. They modulate a coefficient but do not supply its missing scale. The same point applies when a small angle is used as an off-diagonal entry: an angle is dimensionless, while a Hamiltonian entry carries energy.

Check units before integrating the evolution. Otherwise a numerical routine can evaluate a formally typed array while the resulting duration and frequency have no consistent physical meaning.

$$H_{ab} = \varepsilon_{ab}(M_{ab} + M_{ab}^{\dagger})$$

Sources: S04, S05, S07, S09

A consistent static limit

For the source proper-time compensation template, define x=Phi/c². Multiplying its inverse locking factor by the weak-field proper-time factor gives the expression below. The expansion is a first-order statement.

$$\frac{\sqrt{1+2x}}{1+\kappa_i x} = 1 + (1-\kappa_i)x + O(x^2)$$

The choice kappa_i=1 cancels this particular first-order dependence. It does not make the exact expression constant at every order, and it does not derive a nonzero exception for a selected sector. Such an exception needs a physical operator and a composition-dependent observation model.

Keep this comparison distinct from the covariant connection law. A coordinate redshift, a local interaction coefficient and a differential violation are different quantities. Their relation must be calculated in the complete effective action.

Continue with: M12 / M31 / M38 / M43


Sources: S04, S05, S07, S09

Why the variant must be named

The source history contains a polynomial-plus-entropy potential and a tilted double-well potential. Both can be studied mathematically, but they are different functions. Their stationary points, endpoint behaviour and transition structure cannot be combined as though one derivation applies unchanged to both.

The entropy variant uses a linear locking coefficient, a quadratic ordering term, a quartic coefficient and a temperature-dependent entropy contribution. The double well uses a quartic polynomial vanishing at two endpoints before a tilt is introduced.

The manual presents each with its own coefficients and scope. A final canonical action may select one or define a relationship between them. Until then, calculations must name the chosen variant.



Symmetric double-well shape (unscaled)

Sources: S04, S05, S07, S09

The entropy variant

Write K_a for T_eff times d_a. This abbreviation makes the derivatives easier to inspect. The potential includes the mixing entropy alpha log alpha plus one minus alpha times log one minus alpha. Its interior derivatives are singular as either endpoint is approached at positive K_a.

All coefficients in this spacetime potential must carry compatible energy-density units when alpha is dimensionless. A locking energy from the internal Hamiltonian is not the same coefficient without a specified conversion. The kinetic normalisation is treated separately.

The logarithmic function is retained when analysing stability. Replacing it with a local polynomial approximation must state the expansion point and validity range.

$$V = -\mu\alpha + \tfrac{1}{2}(K - r)\alpha^2 + \tfrac{1}{4}u\alpha^4$$

$$V_{\mathrm{ent}} = K[\alpha \ln \alpha + (1 - \alpha)\ln(1 - \alpha)]$$

Sources: S04, S05, S07, S09

The tilted double well

The polynomial variant is lambda times alpha squared times one minus alpha squared in the factorised sense alpha squared times (alpha minus one) squared, plus a linear tilt. At zero tilt, the endpoints are equal minima and the midpoint is a barrier for positive lambda.

Differentiation gives a cubic stationary equation. A nonzero tilt changes the energies and locations of stationary points. It cannot be assumed to preserve both endpoint minima exactly. The field's constrained interval also affects how endpoint configurations are interpreted.

The diagram shows the symmetric dimensionless shape normalised to its maximum. It is a graph of the stated polynomial, not a computed cosmological history.

$$V_{\mathrm{dw}} = \lambda \alpha^2 (\alpha - 1)^2 + \varepsilon \alpha$$

$$V'_{\mathrm{dw}} = 2\lambda \alpha (\alpha - 1)(2\alpha - 1) + \varepsilon$$

Sources: S04, S05, S07, S09

Selecting a calculational model

Specify the potential, kinetic term, sector couplings and allowed field domain together. Then calculate stationary points and fluctuation masses. To claim a first-order transition, compare distinct minima and evaluate the transition conditions; a coefficient sign alone does not complete that analysis.

Bubble nucleation additionally needs a gradient term, dimensional context and an appropriate action. Thermal and vacuum tunnelling calculations use different bounce constructions. The same expression cannot be carried between them without its regime being stated.

The following modules carry out exact local derivatives of the entropy variant and inspect the proposed Casimir inequality. These checks are finite parts of the activation programme.

Continue with: M23 / M24 / M33

Sources: S04, S05, S07, S09

Stability starts with derivatives

A stationary point of an unconstrained interior activation potential satisfies V prime equal to zero. A positive second derivative gives local stability for a single canonically normalised scalar in the absence of coupled directions. Multiple fields require a Hessian and a kinetic matrix.

For the entropy variant, keep the logarithmic term when differentiating. Its derivative is the log of alpha divided by one minus alpha. Its second derivative is one over alpha times one minus alpha. These contributions remain relevant at positive temperature.

The calculation on these pages establishes the local shape of the written potential. It does not determine its coefficients or select a cosmological solution without initial conditions and the coupled field equations.

$$V'(\alpha_*) = 0$$

Sources: S04, S05, S07, S09

Exact interior derivatives

With K equal to T_eff d_a, the first derivative contains the linear locking contribution, the quadratic ordering contribution, the cubic quartic contribution and the entropy logarithm. The second derivative contains K minus r, three u alpha squared and the positive entropy curvature.

For positive K, the endpoint derivatives are not finite. A statement that the minimum occurs exactly at alpha equal to one therefore requires a limiting or constrained-boundary treatment. It cannot be substituted into the interior formula without that explanation.

These exact derivatives provide a direct inspection test for shorthand threshold claims. Any proposed instability criterion must be consistent with the actual curvature of the selected potential.

$$V' = -\mu + (K - r)\alpha + u\alpha^3$$

$$V'_{\mathrm{ent}} = K \ln \frac{\alpha}{1-\alpha}$$

$$V'' = K - r + 3u\alpha^2 + \frac{K}{\alpha(1-\alpha)}$$

Sources: S04, S05, S07, S09

A zero-temperature check

At K equal to zero, the entropy term drops out of this potential. If the unconstrained polynomial is stipulated to have a stationary point at alpha equal to one, its first derivative gives u equal to mu plus r. Substituting this relation into the second derivative gives three mu plus two r.

This result is a potential curvature. It is not automatically a mass squared until the kinetic normalisation is supplied. For a constant coefficient f squared, the corresponding fluctuation mass squared divides that curvature by f squared.

This exact calculation replaces the unsupported shorthand that the same potential always gives mass squared equal to two times its linear locking coefficient.

$$u = \mu + r \quad \Longrightarrow \quad V''(1) = 3\mu + 2r$$

$$m_\phi^2 = \frac{3\mu + 2r}{f^2}$$

Sources: S04, S05, S07, S09

Coupled fluctuations

For several activation fields, evaluate the Hessian at a stationary configuration. The kinetic matrix determines the canonical normalisation. The physical small-fluctuation mass-squared eigenvalues come from the normalised Hessian, not from its raw entries alone.

Mixing with the Higgs, metric or other fields can alter this calculation. A non-minimal gravitational coupling may require a frame transformation and additional field normalisation before a particle mass is identified. The variables and approximations should be explicit.

The mass programme therefore needs both the potential coefficients and the kinetic structure. The manual's testing section records those inputs as prerequisites for a numerical prediction.

Continue with: F04 / M24 / M41

Sources: S04, S05, S07, S09

The proposed filter

The source proposes an ordering drive proportional to a quadratic Casimir and compares it with an entropic cost proportional to sector dimension and effective temperature. This is intended to organise activation thresholds. Its physical force depends on the precise potential and coefficient definitions.

The original inequality is written as r_a greater than four T_eff d_a. If r_a equals B times C2(a), where B collects the remaining factors, the equivalent inequality retains the temperature. Removing it requires a stated rescaling.

A list of dimension-to-Casimir ratios can organise candidate inequalities. It cannot determine the current active gauge group without a fixed cutoff, a valid sector definition and a dynamical solution.

$$r_a = BC_2(a), \qquad r_a > 4T_{\text{eff}} d_a$$

$$B > 4T_{\text{eff}} \mathcal{R}_a$$

Sources: S04, S05, S07, S09

The curvature check

For alpha in the open unit interval, alpha times one minus alpha is at most one quarter. Its reciprocal is therefore at least four. With K positive and u nonnegative, the entropy-variant curvature is bounded below by five K minus r.

Consequently r greater than four K does not, by itself, establish a nonconvex potential or a first-order transition. A range remains in which the stated inequality is met but the lower curvature bound is still positive.

A first-order transition needs competing minima and a comparison of their energies, together with the appropriate transition dynamics. The numerical coefficient in a proposed inequality must follow from that analysis.

$$\frac{1}{\alpha(1-\alpha)} \geq 4$$

$$u \geq 0 \quad \Longrightarrow \quad V'' \geq 5K - r$$

Sources: S04, S05, S07, S09

Dormancy requires an interval

To activate a selected set while keeping another set dormant, the model needs a cutoff lying between the relevant thresholds. Merely saying that unwanted ratios are larger does not establish that they lie above the actual cutoff. The selected parameter interval must be calculated.

Conjugate representations have the same quadratic Casimir in a fixed convention. A Casimir-only drive therefore cannot distinguish them by chirality. Any proposed epoch, initial condition or additional operator that provides the distinction must appear in the calculation.

The source's full-adjoint trace definition also yields the same effective Casimir on every projected block. A more selective operator would be a revised definition, not a missing numerical evaluation of that scalar trace.

CHECK: Calculate the cutoff and all relevant sector thresholds in the same normalisation.

Sources: S04, S05, S07, S09

A continuous order parameter needs a closure

The source constrained trace uses a delta function of a projector. For a single projector, its spectral values are zero and one. That expression therefore has discrete support, with weights W0 and W1 fixed by the chosen state or ensemble.

$$Z(\alpha) = W_0\delta(\alpha) + W_1\delta(\alpha - 1)$$

A continuous free energy needs a further coarse-graining: for example a collective average over many selections, or an ensemble constrained by an expectation value. The continuum potential cannot be inferred from the single-projector spectral delta alone.

Once that closure is specified, derive the potential and gradient terms, compare its actual minima and calculate the roll or tunnelling process. This gives the Casimir-filter programme an explicit route from microscopic operator to an epochal activation law.

Continue with: M14 / M19 / M22 / M31

Sources: S04, S05, S07, S09

Z in the whole framework

The author's wider Z language concerns the transition from a fixed interaction configuration to an evolving one. It includes activation, deactivation, state rotation and internal rearrangement. Its role is larger than a single matrix acting on generation labels.

The explicit order-three gate supplies one finite construction within that programme. It gives a checkable internal action. The full Hamiltonian, activation equations and readout specify how such an action participates in physical evolution.

This edition keeps the conceptual programme and the concrete operator together. It does not reduce the former to the latter or promote the latter into a complete cosmological evolution law.

Sources: S01, S07, S09

Three regimes, stated precisely

0Z denotes the selected stable or inactive-mixing regime. It need not make the whole universe static. Other Hamiltonian terms may remain active, and spacetime fields can still evolve.

+Z names an activation or forward-transformation regime under a specified control rule. -Z names deactivation or the reverse transformation under its corresponding rule. The names do not by themselves identify a derivative of activation with a matrix inverse.

An implementation must state what is being controlled: a coupling coefficient, a field trajectory, a finite gate pulse or a readout. That choice determines what an experiment could detect.

0Z | Specified stable configuration or suppressed selected mixing

+Z | Activation or forward transformation under a stated law

-Z | Deactivation or inverse transformation under a stated law

Sources: S01, S07, S09

A control law is part of the model

The sources write generation coefficients as functions of activation. This makes the strength of an internal operation depend on the background state. It does not automatically make the internal operation drive every activation field.

If the gate commutes with a sector projector, its isolated generator cannot change that sector's occupation. Other terms may still do so. The complete mechanism must identify the connecting operators and the field equation.

The key question is therefore how state transformation, availability and effective readout are coupled. The same question applies whether the transformation is continuous mixing or an exact finite cycle.

$$H_Z = H_Z(\{\alpha_a\}), \qquad \dot{P}_a = \tfrac{i}{\hbar}\langle [H, \Pi_a] \rangle$$

Sources: S01, S07, S09

Names that must remain distinct

The letter Z also appears in partition functions, scalar stiffness coefficients and drawn spatial axes. Those uses do not define the same object. The gate is dimensionless; a stiffness has the units required by its field action; a partition function is a statistical or quantum generating object.

The wider programme may connect these structures through explicit maps, but the shared letter cannot supply the map. This edition uses separate subscripts and descriptions where necessary.

Read the following operator modules as finite constructions supporting the wider dynamical claim. Return to the activation and effective-action modules whenever a physical transition is asserted.

Continue with: M26 / M29 / M31 / M32

Sources: S01, S07, S09

VERIFIED

M26 / 1 OF 4

A matrix with an unambiguous direction

This is the concrete generation-block implementation within the wider Z programme. Use the ordered space C3 tensor C16, with one selected internal sixteen-dimensional copy for each generation label.

For column vectors ordered by generations one, two and three, choose the shift whose columns send the first basis vector to the second, the second to the third and the third to the first. This convention fixes the meaning of forward in the manual.

The displayed matrix is off-diagonal in the generation basis. Its entries are ordinary zero and one values on the three-label factor. Replacing each unit entry by the sixteen-dimensional identity gives the full forty-eight-dimensional block action.

The matrix maps a column of block components to the ordered column consisting of the old third, first and second components. The movement of basis labels and the reordered list of components are two descriptions of the same operation.

Forward shift P

$$\begin{vmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{vmatrix}$$

Column vectors: 1 to 2 to 3 to 1

Sources: S01, S07, S09

The three-step identity

Applying P twice gives the reverse shift. Applying it three times gives the identity. Because a permutation matrix has orthonormal columns, its inverse equals its transpose; for this real matrix the transpose also equals its Hermitian adjoint.

The tensor product with the internal identity preserves these properties. Thus Z is unitary on the selected block and satisfies the required order-three identity. It is not the identity operator because it changes each selected generation label.

These statements are finite matrix calculations. They are not consequences of an analogy with a spatial rotation or a change of epoch.

$$P^3 = I_3, \qquad P^{-1} = P^2 = P^\dagger$$

$$Z = P \otimes I_{16}$$

Sources: S01, S07, S09

Checking a basis vector

Apply the matrix to the first generation basis vector. The output is the second basis vector. Apply it again and obtain the third. The third application returns the first. Repeating the same check from either of the other starting labels gives the same cyclic order.

To check the projector relation, map a vector in the first selected block through Z, apply the second projector and compare with projecting before the shift. The actions agree on every vector in the direct sum. Conjugating the first projector therefore produces the second.

The supplied verification script performs these products using exact integer entries, so no floating-point tolerance is needed for this part of the construction.

$$P(1,0,0)^T = (0,1,0)^T$$

$$P(v_1,v_2,v_3)^T = (v_3,v_1,v_2)^T$$

Sources: S01, S07, S09

Retaining the convention

The transpose matrix is equally valid as a reverse convention. Trouble arises only when the displayed matrix and stated projector cycle disagree. The trial edition fixes that disagreement by using this forward matrix consistently.

If another basis ordering is adopted, record the permutation that changes it. Matrix entries and projector labels must transform together. A written arrow such as one to two is not sufficient unless the basis convention is known.

The forward shift supplies the structural Z operation. Its Hamiltonian use requires the further calculations in the second half of the Z section.

Continue with: M26 / M27 / M29

Sources: S01, S07, S09

Fixing the orientation

Use the forward column-vector convention P acting on label one to produce label two, then label three, then label one. Its transpose is the reverse cycle. Choosing either orientation is legitimate, provided the matrix and projector arrows agree.

For the selected generation block, Z is P tensored with the identity on the internal sixteen-dimensional copy. Applying the operator three times gives the identity. The inverse is its second power.

The physical labels do not determine a matrix convention by themselves. Applying the displayed operator to the basis vectors is the exact test.

$$Z^{-1} = Z^{2} = Z^{\dagger}, \qquad Z^{3} = I$$

Sources: S01, S07, S09

A sign is not an inverse

Multiplying Z by minus one does not produce the inverse cycle. Its third power is minus the identity. Likewise, changing the sign of a Hamiltonian reverses its continuous unitary evolution but is not the same operation as substituting a gate matrix.

This matters for the source's -Z vocabulary. A deactivation trajectory, an inverse finite gate and a negative mixing coefficient are distinct constructions. Their relationship must be set by the chosen control law.

The book preserves the regime names while stating which mathematical action is used in each calculation.

$$(-Z)^3 = -I$$

Sources: S01, S07, S09

Diagonalising the cycle

The cyclic shift has eigenvalues one and the two nontrivial cube roots of unity. Fourier eigenvectors diagonalise it. For the stated forward matrix, a normalised eigenvector with eigenvalue lambda has components one, lambda squared and lambda.

In that eigenbasis the gate is a diagonal phase operator. In the generation-label basis it permutes blocks. Both descriptions represent the same operator, but the projectors must be transformed along with the basis.

The earlier diagonal-versus-shift contradiction arose from keeping generation projectors unchanged while changing the gate basis. Transforming both resolves it.

$$\omega = e^{2\pi i/3}, \qquad f_\lambda = \frac{(1, \lambda^2, \lambda)^T}{\sqrt{3}}$$

$$F^\dagger P F = \mathrm{diag}(1, \omega, \omega^2)$$

Sources: S01, S07, S09

What the spectrum does not fix

An order-three unitary constrains eigenvalues and multiplicities. It does not fix a duration, energy scale or particle interpretation. A continuously evolving propagator needs a Hamiltonian.

Nor does every order-three matrix preserve the E8 Lie bracket. If a gate is identified with an algebra automorphism, that identification needs its full representation action or an explicit bracket-preserving construction.

The full-space module supplies two different extensions. Their different multiplicities demonstrate why a selected 48-dimensional block cannot by itself determine the action on all 248 directions.

Continue with: M26 / M28 / M29 / M48

Sources: S01, S07, S09

A selected block is not the whole operator

The three selected sixteen-dimensional copies occupy 48 internal directions. The remaining 200 directions require an action too. One possible extension leaves them fixed. Another uses a generation-SU3 element throughout the E6 times SU3 decomposition.

Both extensions are unitary and have third power equal to the identity. They are not identical operators. Their eigenvalue multiplicities differ, so their Hamiltonian spectra and projected readouts can differ as well.

This is a useful example of rigidity beginning with a fully specified domain. A finite identity is strongest when every part of the operator has been declared.

Sources: S01, S07, S09

The identity-complement extension

On the 48-dimensional block, each of the three eigenvalues occurs sixteen times. Adding an identity action on the remaining 200 directions increases only the multiplicity of eigenvalue one.

The resulting multiplicities are 216, sixteen and sixteen. This is a valid finite linear construction. It is not automatically the adjoint action of the chosen generation-SU3 element on the full E8 algebra.

The construction may be used as a model operator, but its role must remain distinct from the subgroup action considered next.

$$Z_{\text{simple}} = (P \otimes I_{16}) \oplus I_{200}$$

$$\text{mult}(Z_{\text{simple}}) = (216, 16, 16)$$

Sources: S01, S07, S09

The subgroup-action extension

Under E6 times SU3, the adjoint branches into dimensions 78, eight, 81 and 81. The E6 adjoint is fixed by a transformation acting only on generation SU3. The SU3 adjoint has eigenvalue multiplicities two, three and three.

Each 27 times triplet block contributes 27 to each eigenvalue multiplicity. The conjugate block does the same. The complete action therefore has multiplicities 134, 57 and 57.

This action comes from the declared subgroup element and consequently preserves the full algebraic structure. It still needs the physical spectrum and activation assignment to become a particle model.

$$248 = 78 + 8 + 81 + 81$$

$$\text{mult}(Z_{\text{subgroup}}) = (134, 57, 57)$$

Sources: S01, S07, S09

Exact projectors & trace

For an order-three unitary, averaging the identity and its first two powers projects onto the invariant eigenspace. In the subgroup construction that eigenspace has rank 134. The trace of the complete gate equals 77.

These exact quantities distinguish the extensions without any appeal to an image of a lattice. They can be reproduced directly from the block multiplicities.

The source names generation SU3 as global in the intended construction. That is separate from colour SU3. Whether a symmetry is gauged or global is an action-level choice, not a consequence of the matrix determinant alone.

$$E_1 = \frac{I + Z + Z^2}{3}, \qquad \operatorname{Tr} E_1 = 134$$

$$\operatorname{Tr} Z = 77$$

Continue with: M15 / M26 / M29 / M48

Sources: S01, S07, S09

The source Hamiltonian

The two symmetric combinations in the source generation Hamiltonian are identical for an order-three unitary. Its first and second powers therefore do not supply two independent symmetric directions. Their coefficients combine.

On the three labels, Z plus its adjoint is the matrix with zero diagonal and unit off-diagonal entries. Its eigenvalues are two and a doubly degenerate minus one. A sign change of its common coefficient does not distinguish the two neighbouring labels.

This is the exact behaviour of the written generator. It is not a failure of cyclic structure; it specifies the kind of mixing that structure produces.

$$H_{\mathrm{sym}} = \kappa(Z + Z^{\dagger}), \qquad \kappa = \kappa_1 + \kappa_2$$

Sources: S01, S07, S09

Transition probabilities

With constant kappa and an initial state in label one, the probabilities of reaching labels two and three are equal at every time. Their maximum individual value is four ninths. The remaining probability stays in label one.

This calculation distinguishes a symmetric continuous mixing from an exclusive timed forward shift. Calling both a cycle without checking the propagator would hide the difference.

The formulas are dimensionless after introducing s equal to kappa times elapsed time divided by hbar. The value of kappa and the physical clock mapping remain separate inputs.

$$s = \frac{\kappa \tau}{\hbar}$$

$$P_{1 \to 2} = P_{1 \to 3} = \frac{4}{9} \sin^2 \frac{3s}{2}$$

$$P_{1 \to 1} = 1 - \frac{8}{9} \sin^2 \frac{3s}{2}$$

Sources: S01, S07, S09

A separately proposed directed generator

The Hermitian operator i times Z minus its adjoint supplies an independent orientation-sensitive combination. The foundations investigation introduced it as an additional construction; it is not retrospectively attributed to the original symmetric Hamiltonian.

On the three-label block, choose the constant coefficient below for an interval T. Its exponential equals the forward gate exactly. The reverse sign produces the inverse. This is a finite mathematical result, not a calibrated physical gate duration.

The same expression need not act identically on an unspecified complement. The complete operator domain must still be declared.

$$H_{\mathrm{cyc}} = \frac{2\pi\hbar}{3\sqrt{3}\,T}\, i(Z - Z^{\dagger})$$

$$e^{-iH_{\mathrm{cyc}}T/\hbar} = Z$$

Sources: S01, S07, S09

Connecting the gate to activation

A general Hermitian gate sector can contain symmetric and orientation-sensitive terms, with coefficients controlled by activation. The resulting evolution is time ordered when the complete generators at different times fail to commute.

The finite pulse demonstrates what is required for a directed transformation. It does not derive the coefficient from a Higgs background or a cosmological epoch. That derivation belongs to the broader coupling problem.

This preserves the author's intended motion beyond fixed assignments while replacing a mistaken directional claim about the symmetric generator with the exact calculation.

$$H_Z = \kappa(Z + Z^\dagger) + \chi\, i(Z - Z^\dagger)$$

Continue with: M25 / M30 / M31 / M44

Sources: S01, S07, S09

PROPOSED

M30 / 1 OF 4

A spectrum is not yet a mass hierarchy

The symmetric generation matrix has one eigenvalue different from a doubly degenerate pair. On the selected forty-eight-dimensional block, the multiplicities are sixteen and thirty-two. This spectrum is exact for the stated matrix.

Interpreting it as physical fermion masses requires the effective mass terms and canonical field normalisation. For Dirac fermions, physical masses are singular values of the mass matrix. Signed eigenvalues of a Hermitian internal operator are not automatically those singular values.

The source proposes that Higgs and Yukawa contributions complete the mass structure. Those matrices and their activation dependence must be specified before the observed hierarchy or mixing angles can be derived.



Sources: S01, S07, S09

The fermion calculation

Let M_f be an effective Dirac mass matrix. Left and right unitary transformations bring it to a diagonal matrix of nonnegative masses. In the Standard Model convention, an effective Yukawa matrix produces the mass matrix through the Higgs vacuum scale divided by the square root of two.

The up-type and down-type left diagonalising matrices determine the quark mixing matrix. Neutrino mixing additionally depends on the chosen neutrino mass mechanism. A common internal generation operator does not by itself determine all these matrices.

The model must explain which terms break the cyclic symmetry, lift degeneracies and generate the relevant complex phases. Those are physical completion tasks, not properties of the order-three identity alone.

$$U_L^\dagger M_f U_R = \mathrm{diag}(m_1, m_2, m_3)$$

$$M_f = \frac{v}{\sqrt{2}} Y_f$$

Sources: S01, S07, S09

The common-basis limitation

If two Hermitian mass constructions are functions of the same symmetric generation matrix and the identity, they commute. They can be diagonalised in a common basis. This does not force a measured nontrivial CKM matrix.

Degenerate subspaces permit arbitrary basis rotations whose apparent angles need not represent physical mixing between distinct masses. A realistic hierarchy and mixing analysis must first resolve the degeneracy and identify the charged-current couplings.

The CKM matrix is calculated from the mismatch of the left-handed diagonalising bases, not by simply reading an angle from a spatial cell drawing. The correspondence would require a defined Yukawa construction.

$$V_{\mathrm{CKM}} = U_{uL}^{\dagger} U_{dL}$$

$$[f(A), g(A)] = 0$$

Sources: S01, S07, S09

What a completed derivation would contain

Supply the effective up, down, charged-lepton and neutrino matrices, their kinetic normalisations and the source of their coefficients. State which quantities are inputs, which are fitted and which are calculated from internal geometry or activation.

Then calculate singular values, mixing matrices and phase invariants at a specified scale. Compare with observations using consistent running and uncertainties. A geometric proposal becomes a mass prediction through this complete sequence.

The Z-gate supplies a calculable candidate mixing structure. The trial manual gives that structure a precise scope while recording the extra equations required for its mass interpretation.

Continue with: M29 / M41 / M12

Sources: S01, S07, S09

The layers in one functional

The source master functional brings gravity, activation and matter into one effective description. This is the intended point of contact between the keychain rule and an observable world. Gauge kinetic terms, constraints and auxiliary degrees of freedom must also be present wherever the chosen fields require them.

At this stage the functional is written on an effective spacetime manifold. It organises the local theory once that readout exists. The deeper emergence claim requires the map that supplies this effective description from the interaction substrate.

The functional is therefore a bridge specification. Each contribution has to be explicit before the complete model can generate observables.

Sources: S01, S04, S06, S09

State the action convention

This edition uses Lorentzian signature minus plus plus plus for the displayed effective scalar action. Its quantum weight is the exponential of i times S over hbar. A Euclidean tunnelling calculation instead uses the exponential of minus the Euclidean action over hbar.

Mixing the Lorentzian measure with the Euclidean weight without declaring continuation obscures the calculation. The two uses are retained with their conventions identified.

The total symbol below lists contributions. It does not claim that an unspecified gauge or auxiliary action has already been uniquely derived.

$$S_{\mathrm{eff}} = S_g + S_\alpha + S_{\mathrm{YM}} + S_\psi + S_{\mathrm{aux}}$$

$$\mathcal{Z}_L = \int \mathcal{D}\Phi \, e^{iS_{\mathrm{eff}}/\hbar}$$

$$\mathcal{Z}_E = \int \mathcal{D}\Phi_E \, e^{-S_E/\hbar}$$

Sources: S01, S04, S06, S09

Gauge consistency with varying activation

A spacetime-dependent coupling requires a consistent gauge convention. If the connection is written as the product of a coupling and a gauge potential, its transformation law must account for that product. One cannot vary the coupling while leaving incompatible definitions of the connection and curvature unchanged.

An equivalent description can place a scalar-dependent function in front of a gauge kinetic term while treating the connection convention consistently. The effective physical coupling is then obtained after the appropriate normalisation.

Species permissions must also respect the chosen representation. Arbitrary independent weights cannot replace the generator relations required by nonabelian gauge symmetry.

$$\mathcal{A}_\mu = g(x) A_\mu$$

$$\mathcal{L}_{\mathrm{YM}} = -\tfrac{1}{4} I(\alpha)\,\mathrm{Tr}(\mathcal{F}_{\mu\nu}\mathcal{F}^{\mu\nu})$$

Sources: S01, S04, S06, S09

M31 / 4 OF 4

Vary, solve & read out

Vary the complete action to obtain the field equations, constraints and stress-energy. Specify the state, boundary data and regime. Solve the resulting system or a declared approximation. Apply the detector readout to the solution.

This is the route from the whole theory to a test. A current-source term, a changing kinetic normalisation and an evolving metric can all contribute to the same measurement. They must be treated together when the claimed effect depends on their interplay.

The later modules supply finite examples of this procedure. Completing it for one physical observable is more consequential than adding another uncalculated constant to a prediction table.

Continue with: M37 / M38 / M43 / M50

Sources: S01, S04, S06, S09

A cycle needs a time scale

An order-three operator specifies a three-step internal cycle. It does not state how long a step takes. A continuous dynamics introduces a generator with energy units and a time parameter. Their ratio through hbar determines the dimensionless evolution phase.

The source's proper-time equation already uses an effective metric to define duration along a timelike worldline. Deriving time from a more primary internal structure requires a separate clock construction. Reusing the same proper-time symbol is not that derivation.

An arrow of time additionally concerns ordering or irreversible behaviour. A reversible finite unitary cycle can be traversed in either direction. A bath, coarse-graining or boundary condition may contribute to an arrow, but each requires an explicit model.

$$U(\tau) = \exp(-iH\tau/\hbar)$$

Sources: S01, S07, S09

Proper time and coordinates

In an effective Lorentzian metric with the stated sign convention, proper time along a timelike trajectory is calculated from the line element. The value depends on the metric and trajectory. It is not the number of internal labels.

A photon follows a null path in the standard effective description, so its own proper-time interval is zero. This does not supply a photon rest-frame clock or imply an infinite signal speed. The spacetime causal structure defines local propagation.

An internal clock proposal needs an observable that changes predictably, a calibration scale and a range over which the readout is unambiguous. A cyclic phase alone repeats and must be unwrapped or accompanied by another count.

$$c^2 d\tau^2 = -g_{\mu\nu} dx^\mu dx^\nu$$

Sources: S01, S07, S09

The antisymmetric-expectation check

For a unitary Z, the operator Z minus Z dagger is anti-Hermitian. Its expectation in a normalised state is purely imaginary. Consequently its complex argument can take only the sign-dependent imaginary-axis directions when the expectation is nonzero.

This expectation does not provide a smoothly advancing clock angle. When it vanishes, its argument is undefined. Multiplying that argument by a time scale cannot repair the absence of a continuously calibrated readout.

The result identifies a limitation of a proposed clock shorthand. A different relational observable or a fully specified clock subsystem could be investigated, but it would be a new construction with its own verification.

$$\langle Z - Z^{\dagger} \rangle \in i\mathbb{R}$$

Sources: S01, S07, S09

A tangent is not a transition law

A tangent divergence can mark the edge of a chosen coordinate chart. It does not by itself define a physical state change or a continuation across the divergence. The model needs a dynamical equation or a specified hybrid switching rule.

The earlier expression tan[pi(alpha-1/2)] vanishes at alpha=1/2 and diverges at the endpoints zero and one. Its slope does not diverge at the midpoint. An equation using that driver does not automatically keep alpha within the unit interval.

$$\tan[\pi(\alpha - \tfrac{1}{2})] = 0 \quad \alpha = \tfrac{1}{2}$$

A bounded activation control or a threshold can be constructed and analysed, but it is an additional defined mechanism. The finite cyclic gate, a continuous activation field and a clock readout retain their separate roles.

Continue with: M05 / M20 / M25 / M33

Sources: S01, S07, S09

What an epoch means

I, Magnus Randa, propose that an epoch is a persistent configuration of available interactions. Its identity is given by the activation pattern and its readout, rather than by a date alone. A change of epoch is a transition in that pattern.

This connects cosmology to the same keychain rule used for individual particles. A species may retain permission to a channel while the channel becomes suppressed or reactivated. Matter, propagation and effective constants can then change together.

The source history includes activation and deactivation sequences. This edition retains their physical question without assigning a discarded temperature, duration or observational forecast to an uncomputed trajectory.

Sources: S01, S02

A field equation for the history

In the independent-field effective formulation, activation has a kinetic term and a potential. On a homogeneous expanding background with constant kinetic normalisation f squared, its equation includes acceleration, expansion damping and the potential derivative.

Cross-couplings between activations belong in the full effective potential. They can change stationary points, barriers and transition order. Solving fields independently is a declared approximation, not the full theory whenever those couplings matter.

The equation below assumes that the emergent spacetime description already applies. It supplies the history within that effective stage.

$$f_a^2(\ddot{\alpha}_a + 3H_{\mathrm{FRW}}\dot{\alpha}_a) + \partial_a V_{\mathrm{eff}} = 0$$

Sources: S01, S02

Plateaus & transitions

A plateau is a solution remaining near a stable or metastable configuration. Its persistence depends on the potential curvature, initial conditions, damping and any tunnelling rate. A plotted horizontal line does not establish its physical lifetime.

For a finite barrier, a Euclidean bounce calculation can estimate a transition rate under its approximations. The thin-wall expression requires the appropriate tension, vacuum-energy difference and background regime. It cannot be applied unchanged to every activation potential.

The field history must also respect the defined activation range. If ordinary scalar dynamics would leave that range, the formulation needs a constrained coordinate, boundary rule or a different potential.

Sources: S01, S02

A linked cosmological prediction

Once an activation history is solved, use the same history in couplings, mass templates, stress-energy and propagation. The point is to obtain several related readouts from one solution, rather than selecting unrelated functions for each observable.

For example, a common Higgs factor has consequences for mass ratios; a changing electromagnetic coefficient can affect clocks and transport; and rolling activation energy contributes to expansion. These effects should be evaluated together where the model links them.

An epoch proposal becomes a physical calculation when its solution, matching conditions and observable consequences are given. That is the task of the cosmological branch of V.A.S.K.

Continue with: M12 / M24 / M34 / M35

Sources: S01, S02

Dark matter as inaccessible interaction

The author's hidden-channel proposal separates invisibility from absence. An excitation can occupy an interaction sector inaccessible to ordinary electromagnetic detectors while contributing to the combined geometric readout.

Such a sector could be active for its own excitations and effectively inaccessible to ours. A different case is a sector dormant in the present regime. The two cases should not be collapsed into a single word dark.

To obtain a dark-matter model, specify its effective stress-energy, stability, clustering and permitted exchanges. A missing electromagnetic key explains one kind of non-detection; it does not by itself calculate a halo or abundance.

Sources: S01, S03

The deactivation hypothesis

I propose that some expansion effects can be investigated as the relaxation of previously constraining or compressive interaction configurations. In that account, a change in the balance of active contributions can matter as much as the appearance of a new contribution.

The physical calculation must use the total effective stress-energy. Removing a contribution changes both the source and any energy exchanged during the transition. A decrease in a coefficient is not automatically accelerated expansion.

The question is whether a specified evolving interaction configuration produces the required pressure and energy density through the coupled field equations.

$$\frac{\ddot{a}}{a} = -\frac{\rho + 3p}{6F}$$

Sources: S01, S03

The activation stress-energy

For a homogeneous scalar with positive constant kinetic normalisation and a potential, energy density and pressure have opposite signs for the potential contribution. A slowly evolving potential-dominated configuration can therefore have negative pressure.

For several coupled fields, include all kinetic terms, the complete potential and any energy exchange. The acceleration equation displayed here uses a constant gravitational coefficient F and natural units. A varying F requires the additional terms in the back-reaction module.

This supplies an explicit way to assess the author's relaxation interpretation. The sign of the acceleration follows the solution and its stress-energy.

$$\rho_\alpha = \tfrac{1}{2} f^2 \dot{\alpha}^2 + V$$

$$p_\alpha = \tfrac{1}{2} f^2 \dot{\alpha}^2 - V$$

Sources: S01, S03

A corrected redshift inference

A redshift power law must be derived from its stated regime. Under matter domination, nearly constant potential slope, nearly constant potential energy and overdamped roll, the field velocity is proportional to the inverse Hubble rate. Its squared velocity then scales as a cubed.

Under those restricted premises, one plus the equation-of-state parameter scales as a cubed, or one plus redshift to the power minus three. The opposite power does not follow from the same assumptions. Different histories require a fresh solution.

The renewed account therefore retains dynamical dark-sector tests while removing the earlier unearned universal redshift template.

$$\dot{\alpha} \simeq -\frac{V'}{3f^2 H_{\mathrm{FRW}}}$$

$$1 + w_\alpha \propto a^3 = (1+z)^{-3}$$

Continue with: M33 / M38 / M43

Sources: S01, S03

The earlier-light proposal

The source history asks whether photons emitted in a different interaction epoch could survive a later transition and still reach present observers. This follows naturally from the single electromagnetic-key account: turning off an unrelated channel need not, by definition, remove electromagnetic propagation.

I retain this as a physical proposal to investigate. It requires a continuous or matched propagation history through the transition. An earlier source cannot become an observed galaxy merely by being labelled a ghost galaxy.

The complete calculation must connect emission, transport, frequency evolution and the present detector's interaction permissions.

Sources: S01, S03

Transport & survival

Let Gamma describe a photon-removal rate along a specified path, with inverse-time units in the selected parameter. In a simple transport approximation, the survival probability is an exponential of minus the integrated rate.

The rate can include absorption or conversion when those processes exist in the effective model. Scattering changes the angular and temporal distribution and may need a full transport equation rather than a survival factor.

Changes in the electromagnetic kinetic coefficients can also alter mode matching. A history of nonzero activation alone does not prove negligible attenuation.

$$P_{\mathrm{surv}} = \exp\left[-\int_{t_{\mathrm{em}}}^{t_{\mathrm{obs}}} \Gamma_{\gamma}(t)\, dt\right]$$

Sources: S01, S03

Frequency & redshift

Observed redshift is a comparison between emitted and received frequencies interpreted with the source and detector's clocks. If the theory changes those clocks or matter spectra, the inference must use the same activation history consistently.

An apparent maturity, luminosity or mass inferred from a galaxy spectrum is model dependent. It is not a direct measurement of a quark-mass shift. Stellar populations, dust, abundances, lensing and selection effects can affect the same observations.

The proposed epoch signal must therefore be propagated through a source model and an observation model. It needs a signature that distinguishes it from alternative parameter changes.

Sources: S01, S03

A useful observational target

The strongest target is a correlated pattern: an activation history that predicts transport changes together with changes in matter clocks or emission features. Those linked effects can be tested more strongly than an isolated assertion that a distant object looks unexpectedly old.

Specify which features are shifted, how much under the chosen coefficients, and what independent measurements constrain the same model. No telescope image by itself establishes prior-epoch transmission.

This programme keeps photons central to V.A.S.K.'s cosmological interpretation. They are proposed carriers of interaction history, with survival and readout determined by equations.

Continue with: M02 / M06 / M12 / M43

Sources: S01, S03

The asymmetry question

The CP-Higgs proposal uses mixing between matter-preference, antimatter-preference and Higgs-related directions. The intended mechanism is that activation history and unequal mixing can produce an asymmetric readout.

This is an internal-state construction before it is a baryogenesis calculation. Occupation of a labelled direction is not automatically a conserved baryon number or a baryon-to-entropy ratio. The charges, rates and physical state assignment must be supplied.

The calculation should therefore begin with an explicitly Hermitian generator and a declared charge operator. It can then ask how its expectation evolves.

Sources: S01, S02

A minimal three-state generator

Choose diagonal energies Delta, minus Delta and minus mu_H. Couple the two CP-labelled directions to the Higgs-labelled direction with epsilon-plus and epsilon-minus. Hermiticity fixes the conjugate reverse entries.

Every entry has energy units in a Hamiltonian. A dimensionless asymmetry factor can multiply an energy scale, but it cannot be added directly to an energy without that scale.

The coefficients may depend on activation and temperature under a specified law. Their dependence and the initial state must be fixed before integrating the evolution.

$$H_{++} = \Delta, \quad H_{--} = -\Delta, \quad H_{HH} = -\mu_H$$

$$H_{+H} = \varepsilon_+, \quad H_{-H} = \varepsilon_-, \quad H_{Ha} = H_{aH}^*$$

Sources: S01, S02

Charge evolution & freeze-out

For a fixed charge operator B, its expectation changes through the commutator with the Hamiltonian. If that commutator vanishes and there is no other charge exchange, unitary mixing cannot generate a change in the charge expectation.

A realistic abundance calculation must also specify the expanding background, reaction or decay channels, washout and the entropy normalisation. Freeze-out is determined by the relevant rates relative to the background evolution.

These conditions make the CP-Higgs idea calculable. A difference between two occupation probabilities is a useful diagnostic, but its conversion into a physical abundance remains part of the mechanism.

$$\frac{d\langle B\rangle}{d\tau} = \frac{i}{\hbar}\langle [H, B]\rangle$$

Sources: S01, S02

M36 / 4 OF 4

Geometry, phases & flavour

The source history uses angular language for misalignment and CP structure. An angle between state-space directions, a complex phase in a matrix element and an angle between E8 roots are different quantities.

The root calculation supplies a finite set of root-to-root angles. It does not fix a small CP seed or the observed flavour phase merely by reusing an angle symbol. Those quantities require the explicit representation and interaction map.

The renewed draft keeps the asymmetric-mixing programme and removes the earlier claim that a particular seed had already been forced by the smallest E8 root angle.

Continue with: M13 / M29 / M30 / M42


Sources: S01, S02

Internal structure and local fields

The source proposes that low-energy matter and gauge fields arise from the internal construction. A useful effective template contains spacetime spinor fields, gauge connections, couplings and mass terms. The template can organise calculations once those objects and their symmetries are specified.

A finite internal vector supplies representation components but does not automatically supply all field modes, a Fock space or fermionic statistics. The derivation must construct the local field description and identify the regime in which it applies.

The model's activation-dependent coupling can be inserted into a proposed effective action. Gauge invariance then needs to be checked in that complete action, including the transformation of every field and any space-dependent coefficient.



Sources: S01, S04, S06, S09

A Dirac template

The source uses a Dirac-type equation with gauge connections weighted by the effective sector couplings. The gamma matrices encode the spacetime Clifford relation. The matter field's gauge representation determines how the connection acts on its internal components.

An internal Spin10 spinor label does not itself provide a Lorentz spinor. The spacetime spin structure, frame and kinetic operator must be supplied. A gauge-generator projector likewise does not automatically specify the matter representation on which the gauge field acts.

This template is therefore a target effective description. A completed emergence calculation would derive its fields, coefficients and approximations from the more primary construction.

$$(i\gamma^{\mu}D_{\mu} - m_p)\psi_p = 0$$

$$g_{pa}(x) = g_a^{(0)} w_{p,a}\alpha_a(x)$$

Sources: S01, S04, S06, S09

A propagation check

For the usual positive-energy relativistic dispersion, differentiating energy with respect to momentum gives the group velocity c squared p over E. The squared velocity divided by c squared equals one minus the rest-mass contribution divided by energy squared.

With nonnegative invariant mass squared, the local propagation is no faster than c in that effective description. The massless limit reaches c. These statements follow from the dispersion relation and do not count the number of active sector permissions.

An alternative dispersion or refractive model must be stated and tested separately. Inactive coupling factors do not remove a kinetic term or imply an infinite group velocity by themselves.

$$v = \frac{dE}{dp} = \frac{c^2 p}{E}$$

$$\frac{v^2}{c^2} = 1 - \frac{m^2 c^4}{E^2}$$

Sources: S01, S04, S06, S09

Coarse-graining as a derivation task

Specify the background state, fluctuation variables, retained correlations and field statistics. Expand the chosen action or Hamiltonian, then identify the effective kinetic and interaction terms. State which modes are integrated out and how their effects are represented.

The resulting theory must satisfy the symmetry and conservation requirements appropriate to its fields. A quadratic expansion alone does not prove the full interacting theory or its quantum consistency. It provides a calculable approximation whose scope must be reported.

The public manual can explain this route clearly. The test programme records the action, chiral spectrum and numerical predictions required to complete it.

Continue with: M18 / M38 / M42

Sources: S01, S04, S06, S09

Writing an actual coupled action

A scalar activation can back-react on a metric through its stress-energy. A non-minimal coupling can also make the coefficient of curvature depend on activation. These are familiar mathematical structures in a scalar-tensor effective action.

For one dimensionless activation, write a positive kinetic scale f squared, a potential density V and a curvature coefficient F(alpha). The source's cell-functional proposal can be included by defining F as the reduced Planck coefficient minus xi times that functional, with compatible units.

This gives an explicit effective coupling framework. It does not by itself supply a complete quantum-gravity theory or derive the cell functional from E8.

$$S = \int d^4x \sqrt{-g} \left[\tfrac{1}{2} F R - \tfrac{1}{2} f^2 (\nabla \alpha)^2 - V \right]$$

Sources: S01, S04, S06, S09

The metric and scalar equations

For a constant f and the displayed action, variation of the metric gives the Einstein tensor multiplied by F, the scalar stress-energy and derivative terms in F. Variation of alpha gives its wave operator, the potential derivative and a curvature source involving F prime.

The equations show back-reaction in both directions. The metric affects scalar evolution through covariant derivatives and curvature. The scalar affects the metric through stress-energy and the changing curvature coefficient. A proposed feedback mechanism needs these equations solved together.

If f or the geometric measures also depend on alpha, their derivatives add further terms. The trial equations on this page assume the locally constant kinetic scale stated above.

$$FG_{\mu\nu} = T_{\mu\nu} + \nabla_\mu\nabla_\nu F - g_{\mu\nu}\Box F$$

$$f^2\Box\alpha - V' + \tfrac{1}{2}F'R = 0$$

Sources: S01, S04, S06, S09

Effective strength and stability

In a constant-F background with the conventional minimal matter coupling, the gravitational coefficient is related to G by one over eight pi F in natural units. A changing F requires attention to scalar-mediated effects and the full effective description.

The source's F equal to a Planck coefficient minus xi times the cell functional can make the strength change when activation changes. The direction depends on the sign and field dependence. F must remain in the appropriate positive range for the intended effective gravitational kinetic term.

A stronger gravitational coefficient does not automatically imply accelerated expansion. Acceleration depends on the stress-energy and the complete field equations, including time derivatives of F where present.

$$F = M_{\mathrm{Pl}}^2 - \xi \mathcal{C}_{\mathrm{cell}}$$

$$G_{\mathrm{eff}} = \frac{1}{8\pi F} \quad (F \text{ constant})$$

Sources: S01, S04, S06, S09

What would validate the bridge

Define the geometric functional, its field dependence and coefficient scale. Check units, kinetic signs and the stationary background. Calculate the coupled solutions and their observable consequences. Then compare those consequences with gravitational and cosmological constraints.

The metric equation's derivative terms are essential for conservation. Omitting them while claiming a varying curvature coefficient would give an incomplete effective equation. The same care applies to stress-energy from all other fields and the bath.

The bridge is a proposed effective construction with a concrete calculation route. Its physical parameters and emergence from the internal substrate remain obligations of the framework.

Continue with: M23 / M46 / M33


Sources: S01, S04, S06, S09

The source's geometric interpretation

I, Magnus Randa, propose that a black hole is to be understood as a pseudo-local higher-dimensional configuration whose observable spatial appearance is a readout or shadow. In the theory's language, the primary object is an interaction configuration, not a material ball containing an ordinary visible substance called mass.

The physical construction must specify the map between that configuration, an effective metric and the observer's optical image. A horizon, a spatial slice and a photon shadow remain different readouts. The comparison below calculates the nonrotating effective metric case; it does not equate every shadow with a Euclidean projection.

The activation-weighted mass rule and its possible deactivation are treated in M40. A geometric hypersphere label alone fixes neither the mass nor the signal speed.



Schematic projection; not a spacetime metric

Sources: S01, S04, S06, S09

A standard comparison geometry

For the Schwarzschild comparison case, the horizon radius is two G M over c squared. Its spatial area is four pi times that radius squared. These equations define the ordinary effective reference against which a proposed projection can be compared.

An optical black-hole shadow is not simply that horizon surface photographed. In the nonrotating comparison geometry, the limiting photon-orbit structure determines a critical impact parameter. The apparent angular size also depends on distance and observer conditions.

Calling an image a shadow is therefore acceptable as an analogy, but the model's projection map and the observed optical-shadow calculation must remain separate operations.

$$R_s = \frac{2GM}{c^2}, \qquad A_H = 4\pi R_s^2$$

$$ds^2 = -fc^2dt^2 + f^{-1}dr^2 + r^2d\Omega^2$$

$$f(r) = 1 - \frac{R_s}{r}$$

Sources: S01, S04, S06, S09

A null-orbit calculation

In the Schwarzschild comparison metric, the squared impact parameter for a turning point is r squared divided by one minus R_s over r. Its critical extremum occurs at r equal to three halves R_s. Substitution gives the critical impact parameter three square root of three G M over c squared.

This is an exact reference calculation within that specified geometry. It supplies a concrete difference between horizon radius and optical-shadow scale. It does not derive the V.A.S.K. hypersphere's physical projection.

The proposed map would need to reproduce the relevant lensing and dynamical observables or calculate a measurable deviation under stated conditions.

$$b^2 = \frac{r^2}{1 - R_s/r}$$

$$r_{\mathrm{ph}} = \frac{3}{2} R_s, \qquad b_c = \frac{3\sqrt{3}GM}{c^2}$$

Sources: S01, S04, S06, S09

Mass and higher dimensions

Mass is inferred through dynamical and gravitational effects; a black hole need not be a visible material ball for those effects to be measurable. A higher-dimensional label does not imply that its influence is instantaneous or everywhere at once.

If the model defines a projected mass through activations, calculate how that mass enters the effective metric and stress-energy. If activation vanishes, the definition may give zero projected mass, but it does not establish a conserved invisible remnant without the enlarged dynamics.

The following module gives the projected-mass proposal a precise energy-accounting task. It retains the inactive-mass language as a defined interpretation with explicit observable conditions.

Continue with: M47 / M37 / M40

Sources: S01, S04, S06, S09


A defined projected mass

I use inactive mass or suspended mass for the proposed situation in which a mass contribution is suppressed in the effective readout. The source rule writes the black-hole mass as an activation-weighted sum of sector coefficients. If those activations vanish, the defined contribution vanishes; the resulting geometry must still be solved.

This is the intended interpretation, not a conserved hidden-energy formula. An enlarged system must specify the operator or stress-energy that carries exchanged energy. The proposed auxiliary bath gives a place to formulate that accounting.

The auxiliary bath is the proposed place to account for effective exchange and feedback while retaining a larger linear evolution. Its degrees of freedom and coupling must be specified.

$$M_{\mathrm{eff}} = \sum_a \alpha_a M_a^{(0)}$$



Sources: S01, S04, S06, S09

Continuous changes and jumps

For constant sector coefficients, differentiating the projected mass gives the weighted activation derivatives. A discontinuous activation change gives the corresponding finite mass jump. The source names an abrupt deactivation mass change Randa Suspension.

The jump formula defines the change; it does not supply a transition duration, energy flux or causal propagation law. An instantaneous idealisation may be useful in a model, but its regularisation and observational consequences must be calculated.

If accretion, radiative losses or other processes also change the mass, include them consistently in the effective evolution and in the total energy balance. They must not be counted twice through both activation and an independent loss term.

$$\dot{M}_{\mathrm{eff}} = \sum_a \dot{\alpha}_a M_a^{(0)}$$

$$\Delta M_{\mathrm{eff}} = \sum_a \Delta\alpha_a M_a^{(0)}$$

Sources: S01, S04, S06, S09

The enlarged conservation test

For a closed system with a time-independent full Hamiltonian, its expectation value is conserved under unitary evolution. With explicit time dependence, its derivative includes the expectation of the Hamiltonian's explicit derivative. Merely enlarging the state space does not remove that term.

To show transfer into a bath, define H_system, H_bath and H_interaction and calculate their energy derivatives. Their sum must match the full conservation relation. In a spacetime description, the corresponding covariant stress-energy balance must also be shown.

This is the calculational meaning of conserved on the enlarged space. It cannot be replaced by declaring that energy is stored outside the visible projection.

$$H_{\text{total}} = H_S + H_K + H_{SK}$$

$$\frac{d\langle H_{\text{total}}\rangle}{d\tau} = \left\langle \frac{\partial H_{\text{total}}}{\partial \tau} \right\rangle$$

Sources: S01, S04, S06, S09

Gravitational memory as a proposed signal

A permanent strain change is a gravitational-memory observable. A proposed deactivation event needs its spacetime source, outgoing flux, angular distribution and distance-dependent waveform before a memory amplitude can be calculated.

Declaring an auxiliary bath equivalent to gravitational memory does not specify that source. A spherically symmetric mass change alone is not an arbitrary tensor gravitational-wave burst. The emitted or transferred degrees of freedom must be part of the calculation.

The trial manual records the signature as proposed. A completed model would connect activation evolution, energy balance and the detector readout, with finite duration and causal propagation explicitly stated.

Continue with: M20 / M38 / M43

Sources: S01, S04, S06, S09

Where a mass prediction begins

The theory's claim is that effective constants and masses should be derived from interaction structure and its realised configuration. The following checks identify the normalisation and coefficients a completed mass calculation must provide.

A scalar mass calculation needs a potential, a stationary configuration and a canonically normalised fluctuation. A fermion mass calculation needs an effective mass matrix and its singular values. Both require a scale and specified coefficients.

The source proposes geometric and bath-based ways to determine those coefficients. Their derivations must retain operator ordering, units and the distinction between a potential density and a Hamiltonian energy. A formula made to reproduce an observed mass is a calibration unless its coefficients were independently fixed.

This module gives two finite checks useful for that programme: the canonical scalar relation and a countercheck on a Gaussian matrix shortcut.

$$m_\phi^2 = \frac{V''(\alpha_*)}{f^2}$$

Sources: S01, S02

The standard Higgs comparison

For a canonically normalised Standard Model Higgs doublet with its usual quartic potential, the tree-level relation is mass squared equal to two lambda_H times v squared. This supplies a comparison relation; it does not determine lambda_H from V.A.S.K. geometry.

To use a proposed activation potential for this purpose, specify the map between the activation fluctuation and the physical Higgs field. Include mixing and normalisation. A dimensionless geometric coefficient can enter only through a stated relation that has been derived or defined.

Renormalised masses and couplings also require a scheme and scale. Running a parameter means calculating its scale dependence in a specified effective theory, not changing it to match a target number.

$$m_H^2 = 2\lambda_H v^2$$

Sources: S01, S02

A Gaussian matrix countercheck

Let H equal x sigma_x plus y sigma_z, with independent centred Gaussian variables of variance sigma squared. The Pauli anticommutation relation gives H squared equal to x squared plus y squared times the identity.

The averaged second moment is two sigma squared. The fourth moment is eight sigma to the fourth. Expanding the averaged exponential therefore gives a fourth-order coefficient sigma to the fourth over three. Exponentiating only the averaged second-order term would instead give one half.

The mismatch shows that a scalar Gaussian exponential shortcut is not generally valid for noncommuting random matrices. It does not prevent a properly ordered calculation of the proposed bath reduction.

$$H^2 = (x^2 + y^2)I$$

$$\langle H^4 \rangle = 8\sigma^4 I$$

$$\langle e^{\beta H} \rangle = I + \beta^2\sigma^2 I + \tfrac{1}{3}\beta^4\sigma^4 I + \cdots$$

Sources: S01, S02

Moments, cumulants and a completed coefficient

For a centred scalar Gaussian, the fourth moment is three times the squared variance, while the fourth cumulant is zero. They are different quantities. A derivation that invokes a fourth cumulant must not insert the fourth moment in its place.

To determine a quartic coefficient, specify the variables, operator ordering, covariance and coarse-graining prescription. Carry the expansion to the required order and retain all terms that contribute. Then connect the coefficient to a canonically normalised effective field.

The trial edition supplies the countercheck and the completion procedure. It does not promote an uncomputed quartic to a derived Higgs mass.

$$\kappa_4 = \langle x^4 \rangle - 3\langle x^2 \rangle^2 = 0$$

Continue with: M23 / M30 / M12

Sources: S01, S02

State the fermion spectrum first

Anomaly calculations concern the effective chiral fields and their gauge representations. A sum of sector projector ranks does not supply that spectrum. The source selects one internal Spin10 spinor for each proposed family; its physical field assignment must be stated before the anomaly trace is evaluated.

For the standard all-left-handed family with a sterile neutrino, the representation list contains the quark doublet, conjugate up and down singlets, lepton doublet, conjugate charged-lepton singlet and sterile singlet. The dimensions total sixteen.

The following checks verify that stipulated family. They do not derive the family from the full E8 construction or establish that all other states are physically dormant.

Q_L | (3,2) | Y=1/6

u^c | (3bar,1) | Y=-2/3

d^c | (3bar,1) | Y=1/3

L_L | (1,2) | Y=-1/2

e^c, nu^c | (1,1) | Y=1, 0

Sources: S01, S08, S09, S10

Nonabelian and mixed coefficients

For the cubic colour anomaly, the quark doublet contributes two triplets, while the two conjugate singlets each contribute one antitriplet. The coefficient therefore vanishes. The weak multiplicity must be counted; counting the doublet as only one triplet gives an incorrect result.

The colour-squared hypercharge coefficient uses the two doublet components and the two singlet hypercharges. The weak-squared hypercharge coefficient uses three colour copies of the quark doublet and one lepton doublet. Both sums vanish, with the common fundamental index factor suppressed.

The listed fractional hypercharges use the convention Q equals T3 plus Y. If Q is instead written T3 plus Y over two, those hypercharge labels must be doubled consistently.

$$A_{SU3^3} = 2 - 1 - 1 = 0$$

$$A_{SU3^2Y} = 2\left(\tfrac{1}{6}\right) - \tfrac{2}{3} + \tfrac{1}{3} = 0$$

$$A_{SU2^2Y} = 3\left(\tfrac{1}{6}\right) - \tfrac{1}{2} = 0$$

Sources: S01, S08, S09, S10

Abelian and gravitational sums

The hypercharge-cubed coefficient sums Y cubed over every left-handed field component, including colour and weak multiplicities. The mixed gravitational-hypercharge coefficient sums Y over those same components. The sterile neutrino contributes zero to both sums.

The exact fractions on this page give zero for the cubic coefficient. The linear sum is one minus two plus one minus one plus one, also zero. These are the abelian checks that must accompany the nonabelian calculation.

There are four weak doublets when colour copies are counted: three quark doublets and one lepton doublet. The even count passes the usual SU2 global-anomaly parity test for this spectrum.

$$A_{Y^3} = \frac{1}{36} - \frac{8}{9} + \frac{1}{9} - \frac{1}{4} + 1 = 0$$

$$A_{\mathrm{grav}^2 Y} = 1 - 2 + 1 - 1 + 1 = 0$$

Sources: S01, S08, S09, S10

VERIFIED

M42 / 4 OF 4

What the Z permutation preserves

A permutation of identical families leaves these gauge representations and anomaly sums unchanged when it acts only on the family label. Each stipulated family already cancels, so three copies cancel without a cancellation between different families.

This is a consistency result for the chosen low-energy content. It does not establish a full E8-to-chiral-spectrum derivation. That requires Lorentz assignments, the treatment of the other blocks, gauge-invariant interactions and the claimed activation selection.

The trial manual records the anomaly calculation as VERIFIED within this narrow scope. The spectrum-emergence and dormancy connections remain PROPOSED and are recorded in the claim ledger.

Continue with: M15 / M16 / M48

Sources: S01, S08, S09, S10

A prediction must meet a measurement

The source's six forecast classes are retained as a test programme: neutrino selection rules, flavour evolution, composition-dependent gravity, epoch-dependent masses, sky correlations and possible new resonances. Earlier numerical placeholders are not promoted into current predictions.

The source lists proposed signals in neutrino physics, coupling variation, cosmology and particle production. Each needs a complete observable calculation and a sensitivity statement. A list of signal names does not establish that an experiment can already falsify the model.

Separate an exact selection rule from a small predicted effect. A null prediction needs a symmetry or operator argument that forbids the process. A small effect needs its amplitude, uncertainty and experimental conditions. Both can be useful, but they are tested differently.

Discarded numerical forecasts are not restored in this trial edition. The test programme retains the observable categories and the calculations required to make them sharp.

Sources: S01, S02

The neutrino selection-rule check

Orthogonal neutrino and antineutrino state labels do not alone forbid an interaction that connects them. An off-diagonal operator can have a nonzero matrix element between orthogonal basis states. The interaction terms and conserved charges determine whether the process is allowed.

To derive an exact null neutrinoless-double-beta-decay result, identify the lepton-number structure and exclude the relevant violating operators under the stated symmetry and field content. Orthogonality of labels is only one property of the state basis.

The model must also specify its neutrino mass mechanism. A mass matrix, effective operator and symmetry argument can supply a real selection rule; the keychain names cannot supply it by themselves.

$$\langle \nu | \bar{\nu} \rangle = 0$$

$\langle \nu | O | \bar{\nu} \rangle$ need not vanish

Sources: S01, S02

Non-observation and sensitivity

A missing signal is a falsifier only when the observation was capable of detecting the predicted effect. For a collider process, the expected event count depends on luminosity, cross-section, branching fraction and efficiency. Backgrounds and statistical fluctuations also affect the decision rule.

For an astronomical test, selection effects and interpretation uncertainties must be included. For a clock test, the relevant sensitivity to the changing parameter must be stated. An unspecified nonzero effect can always lie below a finite instrument's sensitivity.

The source's proposed observables become scientifically useful when those ingredients are fixed. The book's ledger records their present status and the missing calculations.

$$N_{\mathrm{exp}} = \mathcal{L}\,\sigma\,\mathcal{B}\,\varepsilon_{\mathrm{det}}$$

Sources: S01, S02

Independent review and publication

Ask reviewers to inspect a defined claim and its supporting calculation. A matrix identity, spectrum construction or effective-action derivation gives a clear review target. A claim of complete unification requires every bridge relevant to that scope to be supported.

Publication, an ISBN and a checksum establish a record of the edition. They do not independently validate its physics. Likewise, an AI response is not an experimental confirmation or a peer-review decision. The manual can be published as a clearly scoped theoretical proposal.

The final reference material makes the edition usable for that purpose. It supplies symbols, equation locations, calculation statuses, outstanding tasks and source provenance within the same volume.

Continue with: F04 / M48 / M12

Sources: S01, S02

IMPLEMENTED

M44 / 1 OF 4

Run the mathematics that is specified

The finite verification programme accompanying this edition checks projector identities, the cyclic gate, its spectra, transition probabilities, the directed pulse, potential derivatives and the listed anomaly sums. These are actual calculations on explicitly defined objects.

The physical meaning of each calculation remains attached to its premises. Running a matrix exponential does not measure a particle interaction. It verifies the propagator for that matrix and can expose an incorrect interpretation of the equation.

This separates a useful computational result from an attractive interface or an imported programme that has not been run. The source archive's implementation projects are not used as physics evidence.

Sources: S01, S08, S09, S10

The minimum record

Retain the input definitions, basis convention, units, equations, initial state and readout. Record the numerical method when an approximation is used, including tolerances and residuals. Exact arithmetic is preferable for rational identities and finite dimension counts.

For a unitary propagation, check norm and probability sums. For an eigensystem, check residuals and multiplicities. For a field trajectory, check the constraint and any conserved or exchanged quantity appropriate to the action.

The calculation record should make it possible to reproduce a failed assertion as well as a successful one. Corrections then remain part of the lineage.

$$\|Z^3 - I\| = 0, \qquad U^\dagger U = I$$

Sources: S01, S08, S09, S10

The directed-gate check

On the specified three-label block, exponentiating the orientation-sensitive generator for its declared duration gives the cyclic shift. The residual between the computed exponential and the target matrix is checked numerically, while the eigenvalue argument supplies the analytic result.

The symmetric generator is checked separately. Its two outgoing transition probabilities are equal, and their sum with the return probability is one. This directly tests the difference between a mixing and a directed operation.

No elapsed physical duration has been inferred from the numerical verification. T remains the declared pulse parameter until a physical coefficient is derived.

Sources: S01, S08, S09, S10

M44 / 4 OF 4

From a finite check to a physical run

For a physical prediction, add the effective field map and independently fixed coefficients. Solve the activation history, construct the generator and kinetic operator, propagate the state and compute the detector response.

The verification resources contain the mathematics and production checks for this edition. They do not certify every programme or logged claim in the supplied archive. Imported code was treated as source material unless a specific execution is recorded.

The next major result should be one complete physical output with its uncertainty and input provenance, followed by an independent comparison rather than a retrospectively chosen target.

Continue with: M12 / M29 / M43 / M50

Sources: S01, S08, S09, S10

A concrete propagation problem

The one-key versus multi-key question can be investigated with a coupled quadratic model. Take two real mode amplitudes with canonical positive temporal kinetic terms, spatial coefficients c1 squared and c2 squared, and a real symmetric mass-squared matrix.

This is a proposed effective submodel, not a completed microscopic derivation. Its eigenvalues are nevertheless exactly calculable. The calculation shows how propagation depends on both channel kinetics and mixing, rather than on a key count alone.

Natural units are used in the displayed frequency equation. The symbols mu1 and mu2 specify the two diagonal mass scales; delta is the off-diagonal mass-squared entry.

Sources: S01, S08, S09, S10

The mode equation

For a homogeneous background, the two-by-two frequency-squared matrix has diagonal entries a and d and off-diagonal entry delta. The two eigenvalues give the squared frequencies of the normal modes.

The square root is real for real parameters because the matrix is symmetric. Positive temporal kinetic terms do not by themselves guarantee nonnegative squared frequencies: the mass and gradient contributions must also satisfy the stability conditions of the chosen range.

This is an exact calculation in the stated submodel. It introduces no measured numerical speed or mass.

$$a = c_1^2 k^2 + \mu_1^2, \qquad d = c_2^2 k^2 + \mu_2^2$$

$$\omega_{\pm}^2 = \frac{a + d \pm \sqrt{(a-d)^2 + 4\delta^2}}{2}$$

Sources: S01, S08, S09, S10

Limits that can be checked

When delta vanishes, the channels decouple and the two eigenvalues reduce to a and d, with their order chosen by size. If the spatial coefficients agree, both modes have the same high-frequency characteristic speed while their mass eigenvalues can differ.

For unequal spatial coefficients, the high-frequency eigenmodes approach the separate channel characteristics when the finite mixing is subleading. At finite frequency, mixing and the mass matrix change the dispersion and group velocities.

Thus more accessible interactions need not imply a fixed lower maximum speed. The actual matrix and the regime decide the result.

$$\delta = 0 \implies \{\omega_+^2, \omega_-^2\} = \{a, d\}$$

Sources: S01, S08, S09, S10

How activation enters

The framework can place activation dependence in c1, c2, the mass scales or delta, provided the effective action and its symmetries justify that placement. A declared formula then predicts how a changing background moves the dispersion branches.

The detector couples to its permitted linear combination of modes. A source may therefore excite both eigenmodes while a detector is sensitive to only part of the resulting state. Arrival time and conversion probability require that full source-to-detector calculation.

This is the operational bridge for the propagation argument. It preserves the author's question about realised constraints and supplies an equation capable of distinguishing possible answers.

Continue with: M02 / M03 / M05 / M06

Sources: S01, S08, S09, S10

VERIFIED

M46 / 1 OF 4

The role of the construction

PSEUDOCUBES STACK NO VOIDS is an early geometric path in the project. Its current role is a supporting spatial construction within the broader interaction-first account. It does not replace the permission algebra, activation dynamics or a physical readout map.

Use a reference cube of edge L. Join its centre to the six square faces, giving six equal pyramids. Split each face pyramid across a face diagonal into two congruent mirror components. The cube then contains twelve primal tetrahedra.

This exact construction is preserved without substituting equal-edge regular shapes. The author's specified component has three distinct edge classes.

Sources: S01, S08, S09, S10

Exact edge data

Choose representative vertices A at the origin, B at (L,0,0), C at (0,L,0), and O at the cube centre. The three centre-to-corner edges are equal, the two base legs are equal, and the remaining base edge is a face diagonal.

At the source normalisation L equals four, the three classes become two square root three, four and four square root two. Their decimal approximations are reading aids only; the exact radicals define the construction.

The normalisation is dimensionless until a physical scale is assigned. Changing its displayed unit does not establish a minimal SI volume.

$$\left\{\frac{\sqrt{3}L}{2}, \frac{\sqrt{3}L}{2}, \frac{\sqrt{3}L}{2}, L, L, \sqrt{2}L\right\}$$

Sources: S01, S08, S09, S10

Filling & the diamond

Each primal component has volume L cubed divided by twelve. Two make a face pyramid. Two pyramids on opposite sides of a shared square base make the specified diamond, containing four components.

The no-void result comes from the face-to-centre partition: the interiors are disjoint and the pieces cover the cube. Equal total volume alone would not prove that coverage.

The diamond is a square bipyramid with unequal edge classes. It must not be replaced by a regular equal-edge octahedral shape when calculating its measures.

$$V_{\text{primal}} = \frac{L^3}{12}, \quad V_{\text{pyramid}} = \frac{L^3}{6}$$

$$V_{\text{diamond}} = \frac{L^3}{3}, \qquad 12V_{\text{primal}} = L^3$$

Sources: S01, S08, S09, S10

Physical interpretation & dimensional discipline

The author's claim of fundamental admissible volumes is a physical postulate in addition to the finite partition result. A spatial readout must state how those components arise and what makes further physical subdivision inadmissible.

For a proposed area-to-volume locking energy, physical area divided by physical volume already has inverse-length units. Multiplication by hbar c then supplies energy. A further inverse reference length would double-count the scale.

This dimensional correction preserves a possible geometric coupling without pretending that the coupling has already selected an observed particle mass.

$$\mu = \hbar c \frac{A_{\text{phys}}}{V_{\text{phys}}}$$

Continue with: M31 / M41 / M47

Sources: S01, S08, S09, S10

Three different spaces

The E8 roots use eight Euclidean coordinates. The selected internal adjoint block has 248 complex components. Effective spacetime has event coordinates and a causal metric. These are different mathematical structures with different purposes.

A projection from internal root coordinates into a spatial diagram can be written explicitly as a linear map. That supplies a reproducible picture, but it is not automatically the emergence of the metric or the matter field equations.

The distinction lets geometry support the interaction programme without becoming an unrestricted source of masses or clock periods.

Sources: S01, S08, S09, S10

Spheres & observed shadows

A Euclidean three-sphere is a hypersurface in four Euclidean dimensions. Its intrinsic three-dimensional measure is two pi squared times the radius cubed. A slice and a projection of that object are different operations.

The author's black-hole hypersphere proposal uses a higher-dimensional interpretation. To turn it into a physical readout, the signature, map, metric and causal structure must be declared. A fourth Euclidean coordinate cannot silently become a Lorentzian time coordinate.

An optical black-hole shadow also depends on photon trajectories and the observer, not simply on the name of a higher-dimensional shape.

$$\mathcal{V}(S^3) = 2\pi^2 R^3, \qquad r_{\text{slice}}^2 = R^2 - w^2$$

Sources: S01, S08, S09, S10

E8 & Leech packing

The optimal packing results in eight and twenty-four Euclidean dimensions provide a mathematical background for lattice-based proposals. They concern congruent sphere packing. They do not by themselves identify the activation potential or a physical vacuum.

The direct sum of three E8 lattices has rank twenty-four and 720 norm-two roots. The Leech lattice has no norm-two roots in its standard normalisation. They are not the same lattice merely because their ranks agree.

An activation-vacuum connection would need an energy functional and a map from its configurations to the packing problem. That is a separate proposed physical identification.

Sources: S01, S08, S09, S10

M47 / 4 OF 4

A clock needs evolution

Twenty-four dimensions do not define a twenty-four-hour period. A clock requires evolving distinguishable states, a phase or other monotonic readout in a declared range, and a relation to an observer's time.

The internal cyclic operator supplies a finite ordering. A Hamiltonian and energy scale supply an evolution. A readout connects that evolution to a clock. Repeating a dimensional number cannot substitute for those ingredients.

The supporting geometry remains available for explicit constructions. Its place in this edition is to clarify possible readouts and exact invariants while keeping the whole interaction architecture primary.

Continue with: M13 / M28 / M32 / M39

Sources: S01, S08, S09, S10

The upstream E8 proposal

Lisi's work is part of the framework's intellectual lineage. His original E8 proposal includes a connection, curvature and action with dynamics. It should not be described as a theory that literally contains no evolution.

The author's fixed root-assignment or 0Z comparison names a restricted conceptual limit. It does not establish that setting one coefficient to zero reproduces Lisi's entire action. That equivalence would require a field-by-field and action-level map.

This edition preserves the E8 connection while accurately stating what is being compared. Attribution is not a claim that Lisi has endorsed V.A.S.K.

Sources: S01, S08, S09, S10

The author's architectural claim

I, Magnus Randa, claim a different route: permissions and activation are primary, and the observable spectrum is a readout of an evolving interaction state. The aim is not merely to rename a fixed embedding.

The car-and-helicopter analogy expresses that change of route. Its mathematical content must be the changed assumptions, the selected operators and the effective spectrum that results. Time dependence alone does not change the representation content of an unchanged algebraic assignment.

The three routes in the source history are permission-based ontology, dynamical availability and an effective-field construction. Each has its own concrete completion conditions.

Sources: S01, S08, S09, S10

The representation-theoretic obstacle

Distler and Garibaldi examine embeddings into real or complex forms of E8 under stated physical representation requirements. A proposed departure should identify the assumptions it changes and then calculate the physical spectrum under its replacement.

The presence of dormant conjugate states does not, by itself, derive an anomaly-free chiral low-energy theory with the required locality and mass terms. If activation gives those states a gap or suppresses their couplings, that mechanism must be calculated.

The finite anomaly sums for the selected families remain valuable checks. They cannot substitute for the construction of those families from the full substrate.

Sources: S01, S08, S09, S10

What a genuine bypass would show

Supply the internal representation, Lorentz assignments, gauge charges, active spectrum and treatment of unwanted modes. Derive the effective kinetic and interaction terms and verify their symmetries. State precisely which hypothesis of the earlier obstruction no longer applies.

If the new architecture satisfies those requirements, its success will be shown by the resulting theory. It does not depend on declaring an earlier theorem false. A theorem about one construction and a viable different construction can coexist.

This is the strongest defensible form of the author's intended claim: a specified architectural departure, followed through to a spectrum and observables.

Continue with: M15 / M16 / M31 / M42

Sources: S01, S08, S09, S10

M49 / 1 OF 4

Symbols as an index to structure

The source stream compresses names, equations, sector rules and tests into a symbolic sequence. Its useful role is an index to a defined theory, not a substitute for the definitions that make the sequence interpretable.

A symbol can point to an operator, action, source record or experimental test. Its meaning must remain recoverable from the manual. Reusing Z for a gate, a stiffness and a partition function needs explicit separation.

This edition treats the encoded kernel as a compact entry point. The public explanation and mathematical definitions remain complete enough to read without guessing what a glyph was intended to mean.

Sources: S01, S08, S09, S10

CP437 & mathematical text

CP437 specifies a byte-to-character convention. Historical screen fonts can display graphics for byte positions that a text codec treats as controls. A faithful archive must state which convention it uses.

The PDF uses embedded fonts and Unicode mathematical typesetting. A strict CP437 export of the equations requires a reversible escape or transliteration system for unsupported symbols. Silent substitution is not a lossless archive.

Byte-level reversibility and a physical projection are different claims. A successful text round trip establishes encoding fidelity, not a dimension-reducing physical emergence theorem.

Sources: S01, S08, S09, S10

M49 / 3 OF 4

SHA-256 & identity

SHA-256 identifies an exact byte sequence with a 256-bit digest. It can detect changes between a supplied file and a recorded reference. It does not prove the truth of the file, authorship or a publication date by itself.

The revision resources record source identities and the generated manuscript. The final PDF's digest is calculated after it is complete and kept in the external verification record. A digest printed inside a file cannot also be that file's ordinary final self-digest without a separately defined construction.

This use of hashing supports evidence discipline without giving cryptographic arithmetic a role as a physical axiom.

Sources: S01, S08, S09, S10

M49 / 4 OF 4

Keep related projects separate

The author's separate computing and interface projects may draw inspiration from V.A.S.K. They are not imported into the physical theory's axioms or treated as measurements of its performance.

Operational logs can show that software ran, made requests or emitted a message. They do not verify the physical assertions contained in generated text. This matters particularly for the large log archive included in the source history.

The renewed manual therefore records implementation lineage in its provenance while keeping its main subject the interaction-first physical framework, its equations and its proposed readouts.

Continue with: F04 / M44 / M50

Sources: S01, S08, S09, S10

Return to the originating claim

I, Magnus Randa, claim that the observable world should be explained from interaction structure: which channels exist, which permissions excitations possess, which channels are realised, and how the resulting state changes.

KEYCHAIN names the permission architecture. V.A.S.K. names the dynamical activation framework. E8 supplies the selected internal substrate. The Z programme supplies transformations within the larger dynamics. Effective fields and geometry are the proposed readout.

The photon and gluon arguments belong at the centre because they ask how realised channels govern propagation. Multi-key matter, hidden sectors and epochs extend the same question.

Sources: S01, S08, S09, S10

What this edition actually establishes

The finite operator constructions have explicit domains and checked identities. The coupling factorisation has exact conditional drift and ratio consequences. The cyclic Hamiltonian has a calculated transition law, and a separately proposed directed generator has an exact pulse.

The scalar potentials have explicit derivatives. The stipulated fermion spectrum passes the listed anomaly sums. The supporting spatial construction has exact edge and volume relations. None of those results needs a retrospective fit to an observed resonance.

These are usable pieces of a mathematical framework. Their physical assembly requires the explicit maps and independently fixed coefficients recorded in the claim ledger.

Sources: S01, S08, S09, S10

A decisive calculation

Choose one current physical target. Specify the microscopic or effective inputs, solve the activation and state dynamics, derive the kinetic or response operator, and compute the observable through the detector readout.

Retain uncertainty from numerical error, input uncertainty and approximation. Identify which inputs were calibrated and which output was withheld from that calibration. Compare the output with independent data at a declared sensitivity.

A correlated drift test or a well-defined propagation model can make this route concrete. The target should be chosen for calculational completeness and discriminating power, not because a desired numerical value is already available.

Sources: S01, S08, S09, S10

M50 / 4 OF 4

A public statement with a technical backbone

The renewed book presents a strong claim and the means to inspect it. Its register distinguishes the adopted model from its calculated consequences and from the physical bridges still being constructed. That distinction is the mechanism by which the argument can improve.

The 248-page body is an editorial structure for a public technical manual. It does not serve as evidence for the selected adjoint dimension. Its purpose is to make the theory readable, navigable and reproducible.

The next edition should advance when a new calculation changes a specific result or completes a named bridge. The source history, definitions and verified results remain available so that progress does not depend on conversational memory.

Continue with: F01 / M12 / M31 / M43 / M44

Sources: S01, S08, S09, S10

GLOSSARY & SYMBOLS

Glossary / 1 of 8

Working definitions for this edition. Locations identify the modules containing their construction and scope.

Amplitude | A complex state coefficient. Its phase can affect interference. A probability follows only after a state and measurement rule are specified. Location: M09, M18.

Activation | Sector availability. It may be a state-derived occupation or an independent effective field in the named formulation. The closure must be stated. Location: M10, M19.

Adjoint | A representation in which the algebra acts on itself by the bracket. The chosen complex E8 adjoint block has dimension 248. Location: M13, M14.

Arrow of time | A proposed ordering or irreversible behaviour requiring dynamical and boundary information. A reversible order-three operator alone does not establish it. Location: M32.

Auxiliary bath | Additional degrees of freedom introduced to account for effective exchange or feedback. Its Hamiltonian, coupling and reduction rule must be specified. Location: M40.

Availability | Whether a permitted interaction is realised in the local regime; not a second name for the permission itself. Location: M01, M10.

GLOSSARY & SYMBOLS

Glossary / 2 of 8

Working definitions for this edition. Locations identify the modules containing their construction and scope.

Bare coupling | The scale g_a^(0) appearing before the permission and activation factors. It is an input until a separate derivation fixes it. Location: M01, M12.

Basis | An ordered set of independent vectors used to express states and operators. A change of basis transforms every relevant object consistently. Location: M27, M27.

Block | A selected representation subspace in a branching. Its dimension is not necessarily a physical particle count and its Lorentz role must be assigned. Location: M15, M16.

Branching | The decomposition of a representation under a subgroup. The component dimensions add to the original representation dimension. Location: M14-M16.

Canonical normalisation | A field redefinition giving the kinetic term its stated standard coefficient. The scalar mass depends on this normalisation as well as potential curvature. Location: F04, M23.

Characteristic speed | A propagation coefficient obtained from the principal kinetic operator. It is distinct from a count of accessible channels. Location: M06, M45.

GLOSSARY & SYMBOLS

Glossary / 3 of 8

Working definitions for this edition. Locations identify the modules containing their construction and scope.

Cartan directions | The commuting directions organising the root description. E8 has rank eight; these directions combine with the 240 roots to give 248 generators. Location: M13.

Casimir | A representation-dependent invariant operator constructed from generators in a fixed normalisation. The full irreducible adjoint Casimir is scalar. Location: M14, M16.

Causality | The propagation and signal constraints of a specified effective spacetime theory. Internal dimensions or inactive coupling labels do not determine signal speeds alone. Location: M32, M37.

Cell measure | An area, volume or other geometric quantity with stated units and length convention. Its use in an energy formula requires a defined coupling. Location: M46, M46.

Conditional observable | An expectation after a specified filter or selection. Its denominator is the squared norm of that filtering operation when nonzero. Location: M11.

Channel | A specified interaction sector and its dynamics. Photon and gluon examples use EM and Strong channels. Location: M02-M03.

GLOSSARY & SYMBOLS

Glossary / 4 of 8

Working definitions for this edition. Locations identify the modules containing their construction and scope.

Density matrix | A positive unit-trace operator describing a state or ensemble. Closed unitary evolution preserves its trace and spectrum. Location: M09, M18.

Dormancy | A model regime in which selected activations or couplings vanish or remain small. Hidden energy, localisation and observability need separate equations. Location: M19, M40.

Effect | A positive operator bounded by the identity that supplies an outcome probability. It need not be an idempotent projector or determine a unique state update. Location: M08, M11.

Eigenspace | The subspace with a specified eigenvalue. The Z phase eigenspaces are not the same decomposition as the three generation-label subspaces. Location: M27, M28.

Emergence map | The rule connecting the proposed primary description to effective fields or geometry. Its input, target, approximation and dynamics must be stated. Location: F03, F03.

Drift invariant | A ratio or derivative fixed by the permission-times-activation factorisation under declared fixed inputs. Location: M12.

GLOSSARY & SYMBOLS

Glossary / 5 of 8

Working definitions for this edition. Locations identify the modules containing their construction and scope.

Epoch | A proposed long-lived activation regime. A cosmological epoch requires field evolution, background geometry and a transition calculation. Location: M19, M33.

Filter | An operator K used to select or modify a state. Its success norm contains K dagger K; a general filter is not automatically a projector. Location: M08, M11.

Fourier basis | The eigenbasis that diagonalises a cyclic shift. It mixes the generation labels and transforms their projectors with the same basis change. Location: M27.

Generation block | The selected three copies of a sixteen-dimensional internal space. Its dimension is 48; a full Z action also specifies the complement. Location: M26, M28.

Hessian | The matrix of second derivatives of a potential at a stationary configuration. The kinetic matrix must be included to identify physical fluctuation masses. Location: M23.

Gluon | The Strong-key example. Self-interaction is within the same nonabelian channel. Location: M03.

GLOSSARY & SYMBOLS

Glossary / 6 of 8

Working definitions for this edition. Locations identify the modules containing their construction and scope.

Hypersphere | A sphere in a specified higher-dimensional geometry. The Euclidean S3 measure has length-cubed units and is not automatically a Lorentzian horizon area. Location: M47, M39.

IETM-inspired | A modular, navigable manual structure with defined procedures and cross-references. This edition does not claim certification under a technical-documentation standard. Location: Preliminary iv.

Intertwiner | A map relating two specified operator actions. Physical identification may require it also to preserve named gauge or Lorentz structures. Location: M48.

Keychain | The organising permission picture for a species. Its mathematical translation is a weighted combination of sector operators with stated algebraic properties. Location: M01, M08.

Kinetic scale | The coefficient f carrying the scale needed for a dimensionless activation’s kinetic term. It enters a canonically normalised scalar mass. Location: F04, M23.

Photon | The electromagnetic-key example used to investigate propagation and the survival of information across regimes. Location: M02, M35.

GLOSSARY & SYMBOLS

Glossary / 7 of 8

Working definitions for this edition. Locations identify the modules containing their construction and scope.

Leech lattice | The specified rootless even unimodular lattice in 24 Euclidean dimensions. It is distinct from the rank-24 direct sum E8 cubed. Location: M47.

Locking term | A proposed diagonal energy assignment to sectors. Closed diagonal evolution changes phases and does not by itself relax occupations to a minimum. Location: M20, M21.

Mass matrix | An effective field-theory object whose singular values give Dirac masses after canonical normalisation. An internal signed spectrum is not automatically this matrix. Location: M30.

Occupation | The probability P_a obtained from a normalised state and projector. Complete orthogonal occupations sum to one. Location: M09.

Permission | The specified access of a species to a sector. Permission weights enter a coupling definition and are not automatically measured probabilities. Location: M01.

Readout | The map from interaction variables to effective fields, geometry or detector observables. Location: F03, M31.

GLOSSARY & SYMBOLS

Glossary / 8 of 8

Working definitions for this edition. Locations identify the modules containing their construction and scope.

Projection | An operation with a defined domain, target and map. A vector projection, spatial slice and optical black-hole shadow are different constructions. Location: M46, M47, M39.

Randa Suspension | The source’s label for abrupt projected-mass loss when activation changes. A physical mechanism also needs a transition and energy-flux calculation. Location: M40.

Root | A vector in the algebra’s root-space description. E8 has 240 roots in its eight-coordinate standard construction; roots are not all 248 adjoint directions. Location: M13.

Triality | An order-three relationship in a stated algebraic construction. Matching names or orders does not identify particular operators or physical generation assignments. Location: M48.

Z-gate | A defined cyclic internal operation within the wider programme of activation and state transformation. Its full action and Hamiltonian use must be specified. Location: M25-M30.

Whole framework | Permissions, availability, dynamics and emergent readout considered together. Location: F01, M50.

EQUATION REGISTER

Equations / permissions & state

E01 | M07 | Sector projector: Hermitian and idempotent; construct it from a stated basis.

E02 | M08 | Weighted permission: K_p=sum w_pa Pi_a. General weights define a filter, not automatically a projector.

E03 | M09 | Occupation: P_a=Tr(rho Pi_a). A complete orthogonal partition has sum P_a=1.

E04 | M10 | Availability coupling: g_pa=g_a^(0) w_pa alpha_a; state the selected activation closure.

E05 | M11 | Conditioned observable: divide Tr(rho K† O K) by Tr(rho K†K), when nonzero.

E06 | M12 | Fixed-input drift: partial ln g_pa=partial ln alpha_a; same-sector species ratios are invariant.

E07 | M17 | Basis projector: Pi=B(B†B)^(-1)B† for independent columns.

EQUATION REGISTER

Equations / activation & Z

E08 | M18 | Unitary evolution: i hbar dot rho=[H,rho]; trace and spectrum are conserved in a closed system.

E09 | M20 | Occupation evolution: dot P_a=(i/hbar)<[H,Pi_a]>+<dot Pi_a>.

E10 | M21 | Connection: D_tau=partial_tau+Gamma_tau; moving Gamma into H requires -i hbar Gamma.

E11 | M22 | Potential variants: retain the Landau-plus-entropy and tilted-double-well definitions separately.

E12 | M23 | Scalar fluctuation mass: V''/f² for constant kinetic normalisation.

E13 | M24 | Casimir threshold: B>4 T_eff d_a/C2(a) when r_a=B C2(a).

E14 | M26 | Forward gate: Z=P tensor I16, with P e1=e2, P e2=e3, P e3=e1.

E15 | M27 | Inverse and Fourier form: Z^-1=Z²=Z†; transform projectors with the basis.

EQUATION REGISTER

Equations / finite gate & fields

E16 | M28 | Complete-action multiplicities: (216,16,16) for an identity complement; (134,57,57) for the stated subgroup action.

E17 | M29 | Symmetric transitions: $P12=P13=(4/9)\sin^2(3\ \mathrm{kappa}\ \mathrm{tau}/(2\ \mathrm{hbar}))$.

E18 | M29 | Directed pulse: $H_{cyc}=2\ \mathrm{pi}\ \mathrm{hbar}\ i(Z-Z\dagger)/(3\ \mathrm{sqrt}(3)\ T)$, giving $U(T)=Z$.

E19 | M30 | Flavour: $V_{CKM}=U_{uL}\dagger\ U_{dL}$. A common commuting circulant structure does not fix nontrivial mixing.

E20 | M31 | Master functional: Lorentzian weight $\exp(i\ S/\mathrm{hbar})$, or Euclidean $\exp(-S_E/\mathrm{hbar})$ after declared continuation.

E21 | M33 | Homogeneous availability: $f_a^2(\mathrm{alpha_ddot}+3H\ \mathrm{alpha_dot})+\mathrm{partial_a}\ V=0$.

E22 | M34 | Scalar density and pressure: $\rho=\mathrm{kinetic}+V$, $p=\mathrm{kinetic}-V$.

E23 | M35 | Simple survival model: $P_{surv}=\exp(-\mathrm{integral}\ \Gamma_\gamma\ dt)$.

EQUATION REGISTER

Equations / readout & tests

E24 | M36 | Charge evolution: d<B>/d tau=(i/hbar)<[H,B]> for fixed B.

E25 | M38 | Back-reaction: F G_mn=T_mn+nabla_m nabla_n F-g_mn Box F.

E26 | M39 | Nonrotating comparison: $R_s=2GM/c^2$; shadow scale $b_c=3\sqrt{3}\,GM/c^2$.

E27 | M40 | Projected mass: M_eff=sum alpha_a M_a^(0); compute energy exchange separately.

E28 | M42 | All-left-handed family: cubic and mixed anomaly sums vanish with listed multiplicities.

E29 | M43 | Expected counts: luminosity times cross-section, branching fraction and detector efficiency.

E30 | M45 | Coupled modes: $\omega_\pm^2=(a+d\pm\sqrt{(a-d)^2+4\,\delta^2})/2$.

E31 | M46 | Spatial partition: primal $L^3/12$; pyramid $L^3/6$; diamond $L^3/3$.

E32 | M46 | Energy from physical measures: mu=hbar c A_phys/V_phys.

CLAIM LEDGER

Permission operators & readout

DEFINED / Species access

K_p encodes the selected permission pattern. Spin, statistics and representation content belong to the effective species construction. See M01, M07, M37.

VERIFIED / Weighted-projector correction

Fractional weights or overlapping selections need not produce an idempotent projector. The general filter readout uses K†K. See M08, M11.

VERIFIED / Complete occupation constraint

A normalised state and complete orthogonal partition imply sum P_a=1. Independent availability fields have different constraints. See M09, M19.

DEFINED / Two activation closures

Occupation-derived and independent-field variants are both retained. A specific calculation must select one or provide its bridge. See M10, M19.

CLAIM LEDGER

Propagation & hidden channels

DEFINED / Photon / gluon examples

The author assigns direct EM and Strong channel permissions respectively. Nonabelian self-interaction does not add a channel by counting. See M02, M03.

PROPOSED / Realised constraints

Dormant permissions need not constrain propagation as active channels do. Calculate the actual kinetic and detector response. See M05, M06.

VERIFIED / Characteristic calculation

For the stated quadratic template, characteristic squared speeds are generalised eigenvalues of B relative to A. See M06, M45.

PROPOSED / Earlier-light survival

An epoch history needs transport, attenuation, frequency matching and a detector readout. Galaxy appearance is not sufficient evidence alone. See M35.

CLAIM LEDGER

Activation mechanisms

VERIFIED / Locking versus occupation

A fixed commuting locking operator alone leaves occupations unchanged. Relaxation needs additional connecting or reduced dynamics. See M20.

VERIFIED / Exactly empty gated sector

Under alpha=P and exclusively activation-product-gated inflow, an exactly empty sector stays empty. State the seed or extra coupling. See M20.

VERIFIED / Potential-specific threshold

The displayed entropy-plus-Landau potential has its own derivative and curvature; a threshold from another potential cannot be imported. See M22, M23, M24.

PROPOSED / Casimir selection

State algebra, representation, normalisation and cutoff. Conjugate Casimirs do not alone separate conjugate sectors. See M14, M16, M24.

CLAIM LEDGER

Z & flavour

VERIFIED / Cyclic block

The forward column convention, order-three identity, inverse and projector cycle agree. See M26, M27.

VERIFIED / Complete action

The identity complement and subgroup action have different multiplicities and remain separately defined. See M28.

VERIFIED / Mixing versus directed pulse

The source symmetric generator is bidirectional. The additional directed generator is separately proposed and its pulse is checked. See M29.

PROPOSED / Physical flavour map

The gate alone gives no completed Yukawa spectrum. Noncommuting effective mass structure and CP data must be supplied. See M30, M36.

CLAIM LEDGER

Time, geometry & gravity

PROPOSED / Emergent readout

Specify how interaction variables produce effective coordinates, fields and metric. A fibre equation already uses a spacetime readout. See F03, M31.

VERIFIED / Clock shorthand correction

The expectation of Z-Z† is purely imaginary, so its argument is not a smooth clock phase. See M32.

VERIFIED / Supporting geometry

The declared cube-centre partition fills its reference cube. Fundamental physical indivisibility remains the author's additional postulate. See M46.

PROPOSED / Back-reaction

The effective scalar-tensor action has explicit metric and scalar equations. Deriving its coefficients from the primary substrate remains a separate task. See M38.

CLAIM LEDGER

Black holes & energy

PROPOSED / Hypersphere interpretation

The author claims a pseudo-local higher-dimensional configuration and an observable shadow. Supply its causal metric and readout map. See M39.

DEFINED / Inactive mass

The source projected-mass rule specifies a readout contribution. It does not establish a conserved sum of bare sector coefficients. See M40.

VERIFIED / Energy accounting condition

The expectation of a closed total Hamiltonian is constant when its explicit time derivative vanishes. Calculate system-bath exchange. See M40.

PROPOSED / Memory signal

A deactivation event needs an outgoing source and angular distribution before a detector strain can be predicted. See M40.

CLAIM LEDGER

Masses, constants & anomalies

VERIFIED / Canonical scalar mass

A potential curvature must be divided by kinetic normalisation. An unfixed geometric multiplier is an input, not a mass derivation. See M23, M41.

VERIFIED / Stipulated-family anomalies

The stated all-left-handed family cancels the listed local anomalies and has even weak-doublet parity. This is not a spectrum derivation. See M42.

PROPOSED / Neutrino selection rule

Orthogonal state labels alone do not forbid all transition operators. State the conserved charge or dynamical selection rule. See M43.

PROPOSED / Parameter closure

Algebraic selection alone does not fix every coefficient, function or initial condition. A no-free-parameter claim requires those dependencies to close. See M41, M50.

CLAIM LEDGER

Cosmology, evidence & provenance

VERIFIED / Conditional redshift result

Under the stated matter-era slow-roll premises, the displayed drift power is (1+z)^-3; other histories require solving their equations. See M34.

PROPOSED / Observational forecasts

The six classes remain a test programme. Earlier unsupported absolute forecasts have not been reinstated. See M35, M43.

VERIFIED / Integrity is not physical proof

CP437 records text conventions and SHA-256 records byte identity. Neither proves a particle or gravity mechanism. See M49.

IMPLEMENTED / Edition verification

The finite checks and document checks were executed for this build. The separate source inventory records coverage and remaining limits. See M44.

SOURCES & REFERENCES

The source record

S01 / Current author instructions and corrections in this conversation, through 17 October 2026, Asia/Bangkok. These govern the renewal's emphasis, claims and project separation.

S02 / -KV-.pdf, 24 pages. S03 / VASK from KEYCHAIN by Magnus Randa as Magnus Randa Esq._compressed.pdf, 514 pages. The latter's photon/gluon discussion and activation-space account are central conceptual sources.

S04 / VASK_R5_Absolution_Ultimata.docx and ModelCard. S05 / VASK_09_Absolution_Specification and its variants. S06 / The scanned PDF titled Photons & Gluons..., six pages; its visible content concerns the master action and the Lisi/E8 bridge.

These sources include AI-written exchanges as well as authorial positions. An AI endorsement is part of the history, not independent verification. The source inventory preserves original filenames, identities and duplicated versions.

SOURCES & REFERENCES

Lisi: E8 & triality

R01 / A. Garrett Lisi. An Exceptionally Simple Theory of Everything. arXiv:0711.0770 (2007). https://arxiv.org/abs/0711.0770

The original proposal explicitly includes a connection, curvature and action with dynamics. The author's 0Z comparison must not erase that content.

R02 / A. Garrett Lisi. C, P, T, and Triality. arXiv:2407.02497 (2024). https://arxiv.org/abs/2407.02497

R03 / A. Garrett Lisi. Division Algebras, Triality, and Exceptional Magic. Advances in Applied Clifford Algebras 36, 45 (2026); arXiv:2609.12112. https://arxiv.org/abs/2609.12112

These are comparison sources. This edition makes no claim that its selected generation operator has already been identified with every triality action in those works or endorsed by their author.

SOURCES & REFERENCES

Representation & anomaly scope

R04 / Jacques Distler and Skip Garibaldi. There is no "Theory of Everything" inside E8. Communications in Mathematical Physics 298, 419-436 (2010); arXiv:0905.2658. https://arxiv.org/abs/0905.2658

The representation obstruction concerns specified embedding requirements. M48 explains the author's intended architectural departure and the spectrum calculation it still needs.

R05 / Edward Witten. An SU(2) anomaly. Physics Letters B 117 (1982), 324-328. DOI: 10.1016/0370-2693(82)90728-6.

M42 performs the ordinary finite anomaly sums for the stated family and counts four weak doublets including colour multiplicity. That check is conditional on the supplied spectrum and does not by itself build an emergent chiral spectrum.

SOURCES & REFERENCES

Packing results & their role

R06 / Maryna Viazovska. The sphere packing problem in dimension 8. Annals of Mathematics 185 (2017), 991-1015; arXiv:1603.04246.
https://arxiv.org/abs/1603.04246

R07 / Henry Cohn, Abhinav Kumar, Stephen D. Miller, Danylo Radchenko and Maryna Viazovska. The sphere packing problem in dimension 24. Annals of Mathematics 185 (2017), 1017-1033; arXiv:1603.06518.
https://arxiv.org/abs/1603.06518

These results concern Euclidean congruent-sphere packing. The twenty-four-dimensional theorem is attributed to the collaboration. The current arXiv record notes updated ancillary details.

The theory's proposed activation-vacuum interpretation requires a separate energy functional and map. The packing theorems supply mathematical background, not a derivation of the physical readout or a clock period.

SOURCES & REFERENCES

Hashing & CP437

R08 / National Institute of Standards and Technology. Secure Hash Standard, FIPS PUB 180-4 (2015). https://doi.org/10.6028/NIST.FIPS.180-4

The digest identifies exact bytes. It does not certify a calculation or replace a publication record. M49 and the provenance pages state the usage in this edition.

R09 / Unicode Consortium vendor mapping archive. CP437.TXT. https://www.unicode.org/Public/MAPPINGS/VENDORS/MICSFT/PC/CP437.TXT

The mapping distinguishes byte decoding from a historical display font's visible glyphs. Unsupported mathematical characters require an explicit reversible escape convention in a strict CP437 archive.

The public PDF uses embedded fonts and vector mathematical expressions. Its visual symbols are not presented as a strict CP437 byte stream.

SOURCES & REFERENCES

Gravitation comparison

R10 / Karl Schwarzschild. On the gravitational field of a mass point according to Einstein's theory (1916). English translation: arXiv:physics/9905030. https://arxiv.org/abs/physics/9905030

The translation and its foreword are distinct from the original metric construction. M39 uses the conventional nonrotating exterior form and calculates its null-orbit shadow scale; it does not adopt the foreword's interpretation.

R11 / C. Brans and R. H. Dicke. Mach's Principle and a Relativistic Theory of Gravitation. Physical Review 124 (1961), 925-935. https://doi.org/10.1103/PhysRev.124.925

This is scalar-tensor background. M38 varies the particular effective action written in the manual; the historical reference is not a derivation of its interaction-space origin.

SOURCES & REFERENCES

Formal sources & revision lineage

S07 / northnorway.tex, byte-identical in the recovered archive to V.A.S.K.by_Max.tex. It records the factorised coupling, action templates and generation operator. Conflicting variants are named rather than merged.

S08 / KEYCHAIN_VASK_Continuity_Record_2026-10-14.txt. S09 / KEYCHAIN_VASK_Foundations_Investigation_Part_1_2026-10-14.txt. These are earlier reconstruction and calculation records, not original independent experimental evidence.

S10 / The attached 256-page Trial Draft 01. Its embedded editable manuscript and build resources were recovered and revised for this edition.

S11 / The appendix-fix ZIP contains a large operational vask.log. Its complete byte stream was hashed and scanned; it supplies software history, not a new physical derivation. The edition's coverage record states the review limits.

SOURCES & REFERENCES

Reproducibility & publication

R12 / Python Software Foundation. hashlib and codecs documentation. https://docs.python.org/3/library/hashlib.html https://docs.python.org/3/library/codecs.html

The companion revision resources identify the build environment, finite verification code, manuscript and source manifest. Exact inputs and calculation residuals are retained for revision.

A public book, a preprint and a refereed article are different publication routes. This trial PDF makes no submission or acceptance claim. A numerical publication probability has not been calculated from an identified statistical model.

The author's claims and the mathematical results should remain separately identifiable in every route. A final edition needs author review and the actual distributor's current production checks. This review copy does not assign an ISBN or perform publication.

PROVENANCE & REVISION

Encoding & reproducible text

The PDF is a reading and review format. Its public text and equations use embedded Unicode-capable fonts. The editable manuscript preserves the same module order and definitions.

A strict CP437 archive is a separate representation with an explicit escape convention. Line endings, byte-order marks and final newlines are part of byte identity. Original sources are preserved rather than silently normalised.

Source diagrams were not treated as proofs. Where a scanned equation was material to the physical account, it was compared with readable formal variants. OCR remains a locator and may misread mathematical symbols.

Source inventory and finite calculation records accompany the draft as embedded revision resources. They are not additional printed manual pages.

PROVENANCE & REVISION

Equation review & corrections

The renewal reviews the conversation's central formulations against the formal source variants and the supplied draft. It checks the manuscript's finite identities and potential derivatives and records physical completion conditions in the claim ledger.

Corrections include general weighted-filter normalisation, complete occupation constraints, locking versus population transfer, the missing temperature factor in a rewritten threshold, and the difference between symmetric mixing and a directed cyclic gate.

Further corrections concern canonical scalar masses, gauge-action conventions, conditional cosmological drift, neutrino transition selection, energy conservation and the distinction between a root diagram and a physical spacetime map.

Earlier AI statements of a completed parameter-free Higgs prediction, automatic theorem bypass, certain instantaneous ghost propagation or guaranteed confirmation are not used as established results. Their underlying research questions remain in the model programme.

PROVENANCE & REVISION

Coverage of the recovered files

The main source archive contains 195 files and 159 distinct byte versions. Its 31 distinct PDF versions contain 3,480 pages. All files were inventoried and decoded text was indexed; low-text pages were separately processed for review.

The separate appendix-fix archive contains one operational log of 12,080,597,812 uncompressed bytes. The whole stream was read for hashing and bounded keyword inspection without importing it as a theoretical manuscript.

Core formal documents, conceptual photon/gluon passages, the supplied draft and its editable resources were used in the reconstruction. The complete source manifest preserves the wider archive's implementation and duplicate material.

This is not a claim to have independently proved every mathematical assertion embedded in thousands of pages of conversation or software logs. The equation register and claim ledger identify the reviewed core and the conditions of its results.

PROVENANCE & REVISION

Trial Draft 02 / production record

The file contains eight preliminary pages and exactly 248 numbered manual pages, for 256 PDF pages. Its 54 four-page units are followed by 32 pages of glossary, equation register, claims, references and provenance.

The renewed organisation places the whole framework and photon/gluon propagation first. The wider Z programme occupies M25-M30; subsequent time, activation and readout modules continue it. Spatial representation occupies M46-M47.

The layout uses a 4 x 9 inch review trim, embedded fonts, vector equations, bookmarks, page labels and internal links. Editable source and verification records are embedded for the next revision.

The final checks cover page count, page labels, content bounds, links, fonts, excluded implementation vocabulary and finite mathematical results. Visual inspection uses rendered pages and section contact sheets. A final physical print proof remains a later production step.